

**BAŞKENT UNIVERSITY
HEALTH SCIENCE INSTITUTE
PHYSIOLOGY DEPARTMENT
PHYSIOLOGY MASTER'S PROGRAM**

**THE EFFECTS OF HIGH DOSES VITAMIN D ON ANXIETY AND
EXPLORATORY ACTIVITY BEHAVIOURS IN RATS**

BY

ZAHOUR GAMAL EDDN ASMAEIL

MASTER'S THESIS

ANKARA- 2021

**BAŞKENT UNIVERSITY
HEALTH SCIENCE INSTITUTE
PHYSIOLOGY DEPARTMENT
PHYSIOLOGY MASTER'S PROGRAM**

**THE EFFECTS OF HIGH DOSES VITAMIN D ON ANXIETY AND
EXPLORATORY ACTIVITY BEHAVIOURS IN RATS**

BY

ZAHOUR GAMAL EDDN ASMAEIL

MASTER'S THESIS

ANKARA- 2021

BAŞKENT UNIVERSITY
INSTITUTE OF HEALTH SCIENCES

This study, which was prepared by Zahour Gamal Eddn Asmaeil within the framework of the Department of Physiology Master's Program, was accepted as the Master's Thesis by the following jury.

Thesis Defense Date: 10 /11/ 2021

Title of the Thesis: The Effects of High Doses Vitamin D On Anxiety and Exploratory Activity Behaviours in Rats

Thesis Jury Members (Title, Name - Surname, Institution) Signature

**BAŞKENT UNIVERSITY
INSTITUTE OF HEALTH SCIENCES
MASTER'S THESIS STUDY ORIGINALITY REPORT**

Date: 09 / November / 2021

Student Name, Surname: Zahour Gamal Eddn, Asmaeil

Student Number: 21910603

Department: Physiology

Program: Physiology Master's program

Supervisor's Title /Name, Surname: Prof. Dr. Nazan Dolu

Title of Thesis: The Effects of High Doses Vitamin D on Anxiety and Exploratory Activity Behaviours in Rats

According to the originality report, which was taken by the plagiarism detection program named Turnitin by the personal / thesis supervisor on 20/ October/ 2021 on the 43-page part of my Master's thesis study mentioned above, which consists of Introduction, Main Sections and Conclusion Section, the similarity rate of my thesis is 18 %. Applied filters:

1. Excluding bibliography
2. Excluding quotations
3. Excluding text parts that contain less than five (5) words of overlap

I have examined the Başkent University Institute Thesis Study Originality Report Obtaining and Using Procedures and Principles. I hereby declare that my thesis work does not include any plagiarism, that I accept all kinds of legal responsibilities that may arise in case the contrary is determined and that the information given above is correct.

Student Signature

ACKNOWLEDGMENTS

Primarily I would thank God for being able to complete this project with success. I would like to thank my supervisor Prof. Dr. NAZAN DOLU for giving me countless opportunities to develop as a scientist over the course of my master's. She has given me invaluable advice and kept me on the right track.

I would also like to thank my dear husband who has offered considerable time and effort to help me. His enthusiasm has had a profound effect on me and is a primary reason that I have developed a strong desire to pursue successfully completing my thesis.

For my parents, who have supported me in many ways throughout the entirety of my education. I hope that I can become the type of person that both of you are.

I would also like to thank my friends and all workers in the laboratory of (Başkent University), especially Mr. Adam.

Finally, my utmost thanks go to the Ministry of Higher Education in my country, Libya, who did not only provide financial support but have stood by me through the good and tough times.

I would like to emphasize that this work was supported by Başkent University Research Council with (DA20/19) research number and Başkent University's Animal Experiments Ethics Committee with number (20/17).

ÖZET

ZAHOUR ASMAEIL, THE EFFECTS OF HIGH DOSES VITAMIN D ON ANXIETY AND EXPLORATORY ACTIVITY BEHAVIOURS IN RATS, BAŞKENT UNIVERSITY, INSTITUTE OF HEALTH SCIENCES / MASTER'S THESIS 2021

D vitamini, güneş ışığına maruz kalmaya tepki olarak deride üretilen yağda çözünen bir vitamindir. Kalsiyum ve fosfat homeostazının yanı sıra nöromusküler ve immünolojik fonksiyon da dahil olmak üzere çeşitli süreçler için gereklidir. Anksiyete, keşif davranışı ve farklı D vitamini dozları arasındaki ilişkiyi incelemek için çok fazla çalışma yapılmamıştır. Mevcut çalışma, yükseltilmiş artı labirentini kullanarak sıçanlarda farklı D vitamini dozlarının anksiyete ile ilgili davranışlar üzerindeki etkisini değerlendirmeyi amaçlamaktadır. yöntemi ve açık alan testi kullanılarak sıçanların keşif davranışları ve genel aktiviteleri üzerine. Bu çalışmada 8 haftalık 36 adet genç erkek Wistar rat kullanıldı. Sham grubu, D vitamini grubu olarak 3 gruba ayrıldılar. Ad libitum 300 IU Vit D içeren yemle beslendiler. Sham grubu (grup 1) sonda ile normal su aldı (n=12). D vitamini grubu (grup 2) 400 IU D vitamini aldı ve son grup (grup 3) 1000 IU D vitamini (n=12) aldı. İlaçlar gruplara 8 hafta boyunca günde bir kez gavaj yoluyla verildi. 8 hafta sonra, sıçanlar yükseltilmiş artı labirent ve açık alan testlerinde test edildi. Açık alanda Grup 2 ($p<0.02$) ve Grup 3 ($p<0.05$) orta alanda geçirilen süre (STC) değerleri Grup 1'e göre çok daha yüksek bulundu. En düşük STP değeri Grup 2 ve Grup 3'te hemen hemen aynıydı. Yetiştirme sayısı Grup 2'de en yüksekti. Yükseltilmiş artı labirent testi değerlendirildiğinde. Grup 2'de diğer gruplara göre kapalı kolda geçirilen süre (STCA) değeri en düşük, açık kolda geçirilen süre (STOA) değeri en yüksek bulundu. kapalı kol giriş sayısı (ECA) değeri Grup 2'de G3'e göre istatistiksel olarak daha yüksekti ($p=0,05$).

Bulgularımız, 400 IU ve 100 IU D vitamini dozlarının ikisi arasında anlamlı bir fark olmaksızın anksiyojenik etkiler sergilediğini gösterdi. Sonuç olarak D vitamini takviyesinin artması anksiyetede artışa neden olur.

Anahtar Kelimeler: D Vitamini, Anksiyete, Açık alan testi, Sıçan, Yüksek artı labirent.

ABSTRACT

ZAHOUR ASMAEIL, THE EFFECTS OF HIGH DOSES VITAMIN D ON ANXIETY AND EXPLORATORY ACTIVITY BEHAVIOURS IN RATS, BAŞKENT UNIVERSITY, INSTITUTE OF HEALTH SCIENCES / MASTER'S THESIS 2021

Vitamin D is a fat-soluble vitamin produced in the skin in reaction to exposure to sunshine. It is necessary for a variety of processes, including calcium and phosphate homeostasis, as well as neuromuscular and immunological function. Not many studies have been conducted to examine the relationship between anxiety, exploratory behaviour, and different doses of vitamin D. The current study aimed to evaluate the impact of different doses of vitamin D on anxiety-related behaviours in rats using the elevated plus-maze method and on exploratory behaviour and general activity of rats by using the open field test. In this study, 8 weeks old age 36 young male Wistar rats were used. They were divided into 3 groups as sham group and two different doses of vitamin D groups. They were fed with feed containing 300 IU Vit D ad libitum. Sham group (Group 1) took normal water by gavage (n=12). The vitamin D group (Group 2) took 400IU vitamin D, and the last group (Group 3) took 1000IU vitamin D (n=12). The drugs were given once a day for 8 weeks to the groups by gavage. After 8 weeks, the rats were tested in the elevated plus-maze and open field tests. In the open field, spent time in the central area (STC) values of Group 2 ($p<0.02$) and Group 3 ($p<0.05$) were found to be much higher than Group 1. While the STC value of Group 3 was the lowest, the STP value was nearly the same in Group 2 and Group 3. The number of rearing was the highest in Group 2. When the elevated plus-maze test evaluated, there were no statistical differences between groups for the spent time in the closed arm (STCA) and spent time in the open arm (STOA), but in Group 2, STCA value was the lowest and STOA value was the highest from other groups. Entry number of closed arms (ECA) value was statistically higher in Group 2 than Group 3 ($p=0.05$).

Our findings showed that 400IU and 1000IU doses of vitamin D exhibited anxiogenic effects, with no significant difference between the two groups. As a result, an increase in vitamin D supplementation causes an increase in anxiety.

Key Words: Vitamin D, Anxiety, Open field test, Rat, Elevated plus-maze

The Başkent University Research Council supported this research with the DA20/19 research number

CONTENTS

ACKNOWLEDGMENTS.....	I
ÖZET.....	III
ABSTRACT	III
LIST OF TABLES	VII
LIST OF FIGURES	VII
LIST OF ABBREVIATIONS	VIII
1. INTRODUCTION	1
2. GENERAL INFORMATION	3
2.1. Vitamin D.....	3
2.1.1. Forms of vitamin D	3
2.1.2. Sources of vitamin D.....	4
2.1.2.1. Sunlight irradiation	4
2.1.2.2. Dietary sources	4
2.1.2.3. Vitamin D supplementation	4
2.1.3. Vitamin D metabolism.....	5
2.1.4. Regulation of vitamin D metabolism and functions	6
2.1.5 Measurement of vitamin D metabolites	7
2.1.6. What is the meaning of vitamin D deficiency?	7
2.1.7. Toxicity of vitamin D	8
2.1.8. Vitamin D and anxiety.....	9
2.2. Anxiety and Exploratory Behavior	9
2.2.1 Animal models of anxiety	10
2.2.1.1. Models based on exploratory behavior	11
2.2.1.1.1 Plus-maze	11

2.2.1.1.1.1 Elevated plus-maze	11
2.2.1.1.1.2. Elevated zero maze	13
2.2.1.1.1.3. Elevated T-maze test.....	14
3. MATERIALS AND METHODS	16
3.1. Experimental Animal	16
3.2. Open Field Test	18
3.3. Elevated Plus Maze	19
3.4. Statistical Evaluation	20
4. RESULTS	21
5. DISCUSSION	25
6. CONCLUSION	28
REFERENCES.....	29
APPENDICES	
APPENDIX 1: PROJECT APPROVAL	
APPENDIX 2: ETHICAL APPROVAL	

LIST OF TABLES

	page
Table 2. 1 Vitamin D intake recommendations from the Institute of Medicine vs. the Endocrine Society	5
Table 2. 2 The range of plasma 25(OH)D values recommended by the Endocrine Society and the Institute of Medicine.....	8
Table 4. 1 Open Field Test Finding of Groups	21
Table 4. 2 The elevated plus-maze finding of Groups.....	22



LIST OF FIGURES

	page
Figure 2. 1. Chemical structures of vitamin D3 (Right) and vitamin D2 (Left).....	3
Figure 2. 2. photobiosynthesis of vitamin D.....	6
Figure 2. 3. Animal models of normal and state anxiety are classified. Models are divided into two types for ease of understanding: Unconditioned reaction tests and conditioned response tests are also available. is used to denote the tests in this article.....	11
Figure 2. 4. Elevated Plus Maze	12
Figure 2. 5. The elevated zero maze (EZM) is seen from the side (left) and from above (right).	13
Figure 2. 6. Elevated T-maze.....	14
Figure 2. 7. Open field test..	15
Figure 3. 1. Shows the Schedule of the Study	17
Figure 3. 2. Open Field Test Apparatus.....	18
Figure 3. 3. Elevated Plus Maze Apparatus.....	19
Figure 3. 4. Comparison of spent time in center and in peripheral between all groups. .	22
Figure 3. 5. Comparison by percentage of spent time and number of enter in the open arms	23

LIST OF SYMBOLS AND ABBREVIATIONS

EPM	Elevated plus maze
EZM	Elevated zero maze
ETM	Elevated T maze
HPLC	High pressure liquid chromatography
IOM	Institute of Medicine
LC-MS/MS	Isotope dilution liquid chromatography tandem mass spectrometry
MED	Minimum erythematous dose
OFT	Open field test
PTH	Parathyroid hormone
UV	Ultra-violet
UVB	Ultraviolet-B irradiance
VDBP	Vitamin D binding protein
1,24,25(OH) ₃ D	1,24,25 trihydroxy vitamin D
1,25(OH) ₂ D	1,25 dihydroxy vitamin D
25(OH)D	25-hydroxyvitamin D
25 (OH)D ₃	25 hydroxyvitamin D ₃

1. INTRODUCTION

Vitamin D plays a critical part in bone metabolism and calcium (Ca) homeostasis [1]. It is also soluble in fats and is formed in the skin because of direct sunlight exposure. Dietary sources can supply the human body with some of its daily needs. Cod liver oil is another substantial supply of this vitamin. Additionally, it is contained in trace amounts in butter, cream, yolk, liver, and supplements. To increase the bioavailability of vitamin D, In the liver, it must be hydroxylated to 25 (OH) D then converted to 1,25 (OH)₂D [2]. Vitamin D inhibits the release of parathyroid hormone (PTH), while PTH increases the production of vitamin D in the kidneys. PTH also regulates ionized calcium levels. Due to the effects of PTH on the stomach and bone, serum calcium is kept in a tightly controlled range [3].

The activated form of vitamin D (1,25(OH)₂D) maintains phosphorus homeostasis and blood calcium since they are both necessary for neuromuscular functions the mineralization of bones [4]. Recent studies were mentioned a possible intricate relationship between brain growth vitamin D levels [5]. Vitamin D's neuroprotective effect is linked to its effects on neurotrophic factor production and release, prevention of oxidative damage to nervous tissue, intracellular calcium homeostasis, and neuromodulator synthesis [6]. Also, some research suggests a connection between vitamin D deficiency and different diseases, including bone conditions like rickets, osteomalacia, and osteoporosis [7], as well as other conditions such as type 1 and type 2 diabetes mellitus, cancer, obesity [8]. Vitamin D is hazardous in high dosages, and reports of toxicosis have been made on individuals who have taken quantities of vitamin D sporadically (sporadic hypervitaminosis D). Signs are (i) loss of appetite, (ii) nausea, (iii) vomiting, (iv) polyuria, (v) polydipsia, (vi) weariness, (vii) mental state alterations, and (viii) constipation. When hypervitaminosis D is referred to as high levels of 25-hydroxyvitamin D in the blood, it is characterized as high levels of 25-hydroxyvitamin D in the serum, hypercalcemia or hypercalciuria, or both (25[OH]D). High intake of vitamin D over a long period as it could lead to changes in the deposition of calcium in the soft tissues (particularly the heart and kidneys), and in more severe scenarios, death. [9].

There are pieces of evidence that low levels of vitamin D can contribute to anxiety, based on the findings of previous studies [10]. There have only been a few research types that established a connection between high doses of vitamin D and anxiety.

In recent years, it has been confirmed a link between anxiety and vitamin D deficiency because many people are spending less time in the sun [10]. A recent study of emerging risk factors for mood disorders in the general population suggests that vitamin D deficiency may have a significant impact on the overall body health [11].

Elevated plus maze (EPM) and open field (OF) tests are used in the research of anxiety in experimental animals. There is an increasing amount of research that suggests a substantial link between vitamin D insufficiency and emotional disorders such as sadness and anxiety. It has been shown in some of these studies vitamin D deficiency increase anxiety [12] and there are some studies using vitamin D in treatment, claiming that it reduces anxiety in humans [13]

In this study, we aimed to investigate the effects of high doses of vitamin D on anxiety-related behaviors. Their effect on anxiety in rats was investigated by the EPM test. In addition, we aimed to detect the effects of high vitamin D doses on exploratory behavior and overall activity of rats via the OFT used in this research.

2. GENERAL INFORMATION

2.1. Vitamin D:

Vitamin D is considered a steroid hormone that can be obtained through sun exposure, nutrition, or even supplements [14]. It is internationally recognized for its significance in the promotion of bone mineral metabolic health and, because of its antirachitic properties, was historically known as the "sunshine" vitamin. Vitamin D has been researched in recent years for its possible function in cancer prevention, cardiovascular disorders and, other conditions [15].

2.1.1. Forms of Vitamin D:

Vitamin D2 (ergocalciferol) and D3 (cholecalciferol) are the mainly two biological forms of Vitamin D. [16]. Ergocalciferol is a plant-derived form and can be synthesized in plants, yeast, fungi, and mushrooms in response to ultraviolet irradiation [17]. The animal-derived form is cholecalciferol, which can be synthesized non-enzymatically from the precursor 7-dehydrocholesterol (7-DHC). Pre-vitamin D is produced when ultraviolet B (UVB) rays strike the skin, which is an unstable intermediate that must be thermally transformed into vitamin D3 before it can be absorbed by the body (Figure 2.1) [18].




Figure 2. 1. Chemical structures of vitamin D3 (Right) and vitamin D2 (Left) [18]

2.1.2. Sources of Vitamin D:

Sunlight exposure, supplements, and nutrition are the main sources of providing vitamin D for the body.

2.1.2.1. Sunlight Irradiation:

After being exposed to sunlight, solar UVB radiation penetrates the skin and converts 7-dehydrocholesterol to pre-vitamin D₃, which later transforms into vitamin D₃. The amount of UV radiation emitted by the comparable skin length is determined by latitude, altitude, air quality, cloud coverage, daylight hours, and the latitudes of the month of the year. Furthermore, the effects of UVB on the skin are influenced by clothing and the use of sun protection. Vitamin D₃ does not become intoxicated as a result of excessive sun exposure. [18,19]

2.1.2.2. Dietary sources :

Since vitamin D is not present in all foods, diet is not considered the main source of it [20]. It is doubtful that people can receive adequate vitamin D from their diet. Unlike skin synthesis, dietary consumption provides Ergocalciferol and Cholecalciferol. Vitamin D-rich food includes oily fish, eggs, cereals, mushrooms, and powdered milk. However, consumption of these items appears to be modest in poor countries [21].

2.1.2.3. Vitamin D supplementation:

Many low-cost additional forms of vitamin D, such as vitamin D₃ and vitamin D₂, are available over the counter [22] in a variety of licensed drugs and food products, such as calcium supplements and fish oil products with up to 500 IU (12.5 g). 400 IU (10g) multivitamins, as well as oil formulations, are widely available in capsules and tablets with 100,000 IU (2,500g) [23]. Recommended vitamin D intake levels are shown in Table 2.1.

Table 2. 1 Vitamin D intake recommendations from the Institute of Medicine vs. the Endocrine Society [24]

Age group	Recommended Dietary Allowance (RDA) per day	Tolerable Upper Intake Level (UL) per day
Infants 0-6 months	400 IU (10 mcg)*	1000 IU (25 mcg)
Infants 7-12 months	400 IU (10 mcg)*	1500 IU (38 mcg)
Children 1-3 years	600 IU (15 mcg)	2500 IU (63 mcg)
Children 4-8 years	600 IU (15 mcg)	3000 IU (75 mcg)
Children and Adults 9-70 years	600 IU (15 mcg)	4000 IU (100 mcg)
Adults > 70 years	800 IU (20 mcg)	4000 IU (100 mcg)
Pregnancy & Lactation	600 IU (15 mcg)	4000 IU (100 mcg)
* Adequate Intake rather than Recommended Dietary Allowance.		

The suggested quantities of vitamin D3 to be supplied per kg food for rats are 300IU. [25]

2.1.3. Vitamin D Metabolism:

Vitamin D requirements can be provided by endogenously (via sunshine exposure) and exogenously (from diet and supplements). However, it makes no difference how we obtain vitamin D: by skin synthetizations, diet, or supplementation [26]. All vitamin D metabolites has low water solubility; they must be delivered by binding to plasma proteins such as the Vitamin D Binding Protein (VDBP) [27]. Before the inactive form of vitamin D3 is hydroxylated, it has to be attached to the VDBP, which carries inactive D2 and D3 to the liver through the blood and lymph, then the 25-hydroxylase enzyme hydroxylases both vitamin D2 and vitamin D3, causing the formation of 25-hydroxyvitamin D (25[OH]D) [28]. As the most abundant circulating form of vitamin D, 25-hydroxyvitamin D (25-OHD) blood levels are frequently used to estimate total body reserves of vitamin D. It is, however, physiologically inactive and must be hydroxylated in the kidney in order to be converted into the biologically active 1,25-dihydroxyvitamin D (1,25[OH₂] D) (Figure 2.2) [29]. Endocytic receptors, such as megalin and cubilin, can be found in the kidney's proximal tubule cells. They are in charge of VBP reabsorption, which leads to the intracellular conversion of 25(OH)D to 1,25(OH₂) D in the presence of 1 α -hydroxylase [30].




Figure 2. 2. Photobiosynthesis of Vitamin D [14]

2.1.4. Regulation of Vitamin D Metabolism and Functions:

Vitamin D's major function is to keep calcium, phosphorus, and bone homeostasis. Vitamin D is responsible for maintaining adequate and regular calcium levels. Its deficiency has a significant role in the development of bone mineralization illnesses such as osteoporosis and fractures. Low blood calcium levels induce the parathyroid gland to emit PTH, which triggers 1-hydroxylase in the kidneys to produce 1,25(OH)₂D, which elevates calcium levels by three different mechanisms. [31] increases bone mobilization through increased intestinal absorption interacting with PTH to stimulate reabsorption in the kidneys and boosting intestinal absorption although it is suggested that bone cells can convert 25(OH)D to 1,25(OH)₂D when calcium availability is limited. Dietary calcium is usually favored over bone mobilization [32]. A single vitamin D receptor is responsible for vitamin D hormone functions and is found in osteoblasts, enterocytes, and renal tubular cells as target cells. It was recently discovered that tissues and cells that do not regulate calcium and

phosphate metabolism have 1,25(OH)₂D₃ receptors. This involves parathyroid gland cells, the colon, the ovaries, the pituitary gland, promyelocytes, skin keratinocytes, and lymphocytes. This demonstrates that the repercussions of vitamin D insufficiency extend beyond bone metabolism [33]. Vitamin D deficiency is linked to the pathophysiology of different disorders, including hypertension, diabetes mellitus, tumor growth, autoimmune diseases, mood disorders (depression, anxiety, stress) and infections. [34]

2.1.5 Measurement of Vitamin D Metabolites:

One of the most prevalent vitamin D metabolites is 25(OH)D. So, its serum or plasma concentrations are frequently employed to assess vitamin D status. If valid measurements are conducted at room temperature and after up to four freeze-thaw cycles, the concentration of 25(OH)D in serum samples is stable [35]. 25(OH)D concentrations can be determined using immunoassays developed in the early 1980s. To assess the quantity of 25(OH)D, competitive protein binding assay and other methods can all be utilized [36]. LC-MS/MS, in contrast to the majority of commercial immunoassays, is capable of concurrently measuring 25(OH)D₂ and 25(OH)D₃ (total 25(OH)D = 25(OH)D₂ + 25(OH)D₃) and is thus the gold standard method for evaluating vitamin D status [37]. Additionally, LC-MS/MS may be utilized to quantify vitamin D metabolites 24,25(OH)₂D₂ and 24,25(OH)₂D₃ reliably. 1,25(OH)₂D is frequently assessed using an enzyme-linked immunosorbent assay or a chemiluminescent immunoassay, which is especially difficult considering that circulating levels are hundreds of times lower than those of its precursor 25(OH)D [36].

2.1.6. Meaning of Vitamin D Deficiency:

A low level of 25(OH)D in the blood indicates to be a deficiency of vitamin D, as it is widely used in assessing the level of vitamin D in the blood. The explanation for this is that 25(OH)D has a relatively long half-life of around 2-4 weeks. [38]. Furthermore, since the 25-hydroxylation stage is thought to be poorly regulated by metabolism, 25(OH)D should represent vitamin D intake [39]. Low level of 25(OH)D results in vitamin D deficiency. There is no agreement on what constitutes a normal 25(OH)D concentration. [40] There are currently numerous recommendations, it has been proposed that vitamin D status be characterized according to 25(OH)D concentration in humans (Table 2.2) [41] as follows: deficiency (less than 20 ng/ml), insufficiency (21–29 ng/ml), sufficiency (30–100 ng/ml).

Other Institute of Medicine (IOM) guidelines, on the other hand, state that serum 25(OH)D levels between 16 and 20 ng/ml (40 and 50 nmol/l, respectively) are regarded sufficient, and that levels exceeding 50 ng/ml (125 nmol/l) should be evaluated for possible harmful effects on health (Table 2.2.) [42].

Table 2. 2 The range of plasma 25(OH)D values recommended by the Endocrine Society [41].

Vitamin D status	Endocrine Society	Institute of Medicine
Deficient	below 20 ng/ml (50 nmol/l)	below 16 ng/ml (40 nmol/l)
insufficient	21 – 29 ng/ml (52 – 72 nmol/l)	
Sufficient	30 – 100 ng/ml (75 – 250 nmol/l)	16 – 20 ng/ml (40 – 50 nmol/l)

2.1.7. Toxicity of vitamin D:

High vitamin D doses could possibly lead to hypercalcemia. During early stages of vitamin D poisoning, people have gastrointestinal difficulties such as anorexia, diarrhea, constipation, nausea, and vomiting. Furthermore, numerous additional problems, such as aching bones, fatigue, headaches, a racing heartbeat, a decrease in appetite, and muscular and joint discomfort, commonly occur in conjunction with the aforementioned symptoms. Kidney stone illness can frequently be found in those who wake at night to urinate, accompanied by increased thirst, weariness, anxiety, and itching [43]. The quantity of vitamin D produced by one MED is multiplied by 10,000–25,000 IU of oral vitamin D, the amount of UVB radiation required for vitamin D sufficiency may be calculated [44]. The MED refers to the amount of time it takes for the skin to turn pink. The time duration differs

according to the location, skin pigmentation, age, and body fat percentage. It is worth mentioning that sunlight eliminates any additional vitamin D [45].

2.1.8. Vitamin D and Anxiety:

The role of vitamin D is to participate in the processes of development and growth, as well as to maintain the normal structure of the skeleton by regulating calcium metabolism in the body that occurs as a result of increased calcium absorption in the intestine [46]. However, Vitamin D is an active neurosteroid that is thought to modulate a wide range of brain processes. where it is synthesized in the brain and crosses the blood-brain barrier. also, vitamin D receptors have been found on neurons and glia in numerous parts of the human brain, including those involved in emotional regulation. [47] As previously noted, vitamin D's most well-known role is to maintain blood calcium and phosphorus balance, so a rise in vitamin D levels results in hypercalcemia, which causes neuropsychiatric dysfunction. previous research has found the patients with mild cases of hypercalcemia may show different cognitive abnormalities, whereas patients with severe cases of hypercalcemia may present with altered mental state, psychosis, and confusion. [48]

Many research demonstrates a link between hypovitaminosis D and cognitive fundamental and executive role, depression, bipolar disorder, and schizophrenia. Vitamin D enables neuronal receptors in regions involved with behavior control, encourages neurotrophic release and protects the brain with buffering anti-oxidant and anti-amative defenses against vascular injury. [49] Also, vitamin d regulates the release of nerve growth factor (NGF) which is a chemical for the neuronal survival of the hippocampus and cortical neurons [50]. Vitamin D may have autocrine/paracrine characteristics in the human brain, based on the extensive distribution of 1-Alpha-Hydroxylase [51].

2.2. Anxiety and Exploratory Behavior:

Anxiety is a common reaction to stress and is defined as a feeling of worry, uneasiness, or fear about the condition in one's life. Anxiety can present itself in a variety of ways. Anxiety and depression are frequently linked, as both have a destructive effect on one's life. While depression and anxiety are often treatable with medication, the majority of people do not seek treatment for these conditions Individuals suffering from anxiety disorders have a tendency repeat negative thoughts and fears, which prompts them to avoid specific situations out of dread [52]. Anxiety issues affect women more than twice as much as men. Social

phobia disorder, and specific phobia disorder are different forms of anxiety disorders. Anxiety manifests itself physically in a variety of ways, including tension, sweating, elevated heart rate and blood pressure, trembling, dizziness, and rapid heartbeat. [53].

Exploratory behavior, which enables the gathering of information about novel objects, is critical for survival due to the increased opportunities for finding food, water, a mating partner, and shelter that it provides. At any one-time, exploratory behavior is affected by conflicting motivations to investigate a harmful novel area or to remain in safe and familiar settings. So, an animal's behavior in a novel habitat is always impacted by both curiosity and neophobia. Exploration-based behavioral tests in mice are frequently used to assess the anxiety-related effects of psychoactive medicines where it was discovered that animals' exploratory behavior is affected by extreme anxiety. [54].

2.2.1 Animal Models of Anxiety:

Animal models of psychiatric disorders try to show different aspects of the human experience, ranging from behavioral changes that indicate the emotional state to disease etiology and treatment effects. Animal models are "experimental preparations developed in one species for the aim of understanding phenomena occurring in another species," according to McKinney [55]. animal models can aid in the understanding of molecular systems involved in anxiety, as well as the screening and development of new drugs for their treatment that would be unfeasible in humans. The genetic foundation of anxiety has been established through human studies, and animal studies have been utilized to try to better define its genetic determinants. Animal models in the realm of anxiety research can be divided into two categories (i) models based on unconditioned responses; and (ii) models based on conditioned responses. The first category includes models based on rodent exploratory behavior, models based on rodent social behavior or nonhuman primates, and models based on the somatic stress response. Other paradigms that do not neatly fit into any of the other divisions, such as anxiety, are summarized in the fourth group [56].




Figure 2. 3. Animal models of normal and state anxiety are classified. Models are divided into two types for ease of understanding: Unconditioned reaction tests and conditioned response tests are also available. is used to denote the tests in this article. [56]

2.2.1.1. Models Based on Exploratory Behavior:

2.2.1.1.1 Plus-Maze:

2.2.1.1.1.1 Elevated Plus-Maze:

Anxiety animal models such as the elevated plus-maze are unquestionably among the most commonly employed in current preclinical anxiety investigation. The test was developed as a result of the early finding that now in mazes with closed and open alleyways, in enclosed alleys, rats routinely demonstrate higher levels of exploration than open alleyways. However, when given a choice between alley types, rats choose to avoid those that do not have walls around them. This discovery resulted in the first investigations into the possible utility of an elevated 'X'-maze as paradigm of anxiety, which was conducted some 30 years later [57]. These researchers used a straightforward maze configuration that was elevated

two feet above the floor level and comprised two open and enclosed arms. Researchers also verified that rats avoid open arms, but they also demonstrated that diazepam reduced open arm avoidance while picrotoxin increases it [58]. Following this initial report, Comprehensive assessments of the test (that called as the elevated 'plus'-maze) to usage with rodents were conducted out, and more current studies suggest that it may be beneficial for a variety of other species. Beginning with the plus-maze, spatiotemporal measures of anxiety (i.e., the time spent in open arms as a percent or ratio of total arm time, and the frequency of open arm entries as a percent or ratio of total arm entries) have been the primary indices used to assess anxiety, while the total number of entries in all arms have frequently been used to measure general activity [59].



Figure 2. 4. Elevated Plus Maze [60].

2.2.1.1.1.2. Elevated Zero Maze:

It is a recently developed device of the plus-maze. this test, like the elevated-plus maze, is predicated on two opposing inherent tendencies: investigating a new environment and avoiding elevated and open spaces that represent predator risk circumstances. It consists of a circular elevated platform with two open regions opposite each other and two closed regions animals are put in one of the closed regions chosen as the starting region and anxiety-related behaviors are assessed both manually and through a video system [61].




Figure 2. 5. The elevated zero maze (EZM) is seen from the side (left) and from above (right) [62].

2.2.1.1.1.3. Elevated T-maze test:

The elevated T labyrinth is a 3-armed experimental setup made of Plexiglas of the same size (50x12cm) at a length of 50 centimeters above the ground. One of them is closed with 40 cm high side walls made of Plexiglas and is placed perpendicular to two open arms standing opposite each other. T maze with different sizes for mouse and rat is used for rats. The Elevated T-maze takes use of the subjects' natural dread of heights and open spaces. Anxiety in rodents can be assessed by measuring inhibitory avoidance in the face of conflicts, whereas panic can be assessed by conducting one-way escape tests on the Elevated T-maze [63].




Figure 2. 6. Elevated T-maze. [64]

2.2.1.1.2. Open field test:

The Open Field Maze (OFM) was created in 1934 as a test to assess emotionality in rodents. It has become one of the most commonly utilized measurements of behavior in animal psychology. It allows for a simple and quick assessment of well-defined behaviors while requiring minimal training for the test participant and little to no specialist training for the person conducting the test. These characteristics have led to the open field maze's widespread use in studies involving different animal species [65]. Its popularity stems in part from the psychological and physiological benefits it provides. Considered Rodents are nocturnal animals that like to stay in the dark and avoid strong light. This must be considered while doing an open field test, which is a frequent observational strategy. The animal is removed from its cage and placed in an unfamiliar, well-lit arena that allows it to walk about freely [66]. Due to its photophobia, the animal avoids brightly lighted open regions and prefers to remain close to the walls. The region is separated into peripheral and core sections, which may be used to monitor movement and rearing. Exploratory or locomotor behavior is evaluated, as well as an autonomic activity like urination and feces. Infrared beam array technologies are used to track automated movement, rearing, and time spent in specific parts of the open field. Furthermore, comparable to the elevated plus-maze, the open field behavior is highly responsive to a variety of internal and external stimuli. [67].



Figure 2. 7. Open field test [68].

3. MATERIALS AND METHODS

3.1. Experimental Animal:

We used a total of 36 male Wistar rats that were 8 weeks old. Animals were supplied from the Başkent University Experimental Animal Breeding Center. They were placed in normal cages and provided food and water ad libitum. All animal tests were conducted in Başkent University, Experimental Animal Research Center. The tests were conducted with the agreement of Başkent University's Animal Experiments Ethics Committee (20/17).

3.2. Experimental Groups:

Equal numbers of rats were assigned to three groups ($n = 12$).

- Sham group (Group 1) [$n=12$]: Each rat received normal drinking water via oral gavage.
- Vitamin D at a 400IU dose (Group 2) [$n=12$]: Each rat received 400 IU/day of vitamin D via oral gavage [69].
- Vitamin D at a 1000IU dose (Group 3) [$n=12$]: Each rat received 1000 IU/day of vitamin D via oral gavage [70].
- Vitamin which we used in experiment Devit-3 oral drops, each 15 ml drop contains 50,000 IU of vitamin D3 (Deva Holding A.Ş., Turkey). The active ingredient cholecalciferol (vitamin D3) is produced from sheep's wool fat.
- For eight weeks, the groups received the drugs once daily. After this time, the rats are tested. The study's schedule was shown in (Figure 3.1).




Figure 3. 1. The Schedule of the Study

3.3. Open Field Test:

We did the open field test after completing the period of giving the vitamin D [8 weeks] for 2 days in the morning consecutively for each group starting with the sham group in the first 2 days then the low-dose group on the second 2 days and high dose group in the third 2 days. During this test, each rat was placed in the corner of the box (OFT apparatus consists of a wood box ($100 \times 100 \times 36$ cm³) and the test period lasted 10 minutes. A video recording system was used to determine and quantify the overall distance traveled, the number of reared, and the number of defecations. After testing each rat for ten minutes, the device was cleaned with alcohol before began the new test (figure 3.3) [71].



Figure 3. 2. Open Field Test Apparatus

3.4. Elevated Plus Maze:

We did the EPM test after completing the period of giving the vitamin D [8 weeks] for 2 days in the afternoon consecutively for each group starting with the sham group in the first 2 days then the low-dose group on the second 2 days and high dose group in the third 2 days. where the EPM apparatus consists of two open arms (48 cm long, 10 cm broad each) that are perpendicular to two arms open arms that have the same size, that are enclosed by 50-cm high walls. Each open and closed arm is divided into three parts (16×10 cm each) and a Centre open segment that connects the two arms. the rat was first placed in the central segment of the EPM, facing one of the open arms the experiment was started after a 5-minute habituation period. An exploratory activity measurement was recorded for the number of open and closed segments crossed by each rat. The proportion of entries into open arms was computed using the following formula: percent entries= (entries into open arms/total number of entries $\times 100$). Similarly, the percentage of time spent in open arms was calculated as follows: percent time= (time in open arms/ (time in open arms + time in closed arms) $\times 100$) [72]. After each rat was tested for 5 minutes, alcohol was used to clean the apparatus.




Figure 3. 3. Elevated Plus Maze Apparatus.

3.5. Statistical Evaluation:

The data is evaluated using descriptive statistics and mean comparisons between all study groups. One-way analysis of variance (ANOVA) was used to examine anxiety and exploratory activity, followed by Tukey's multiple ranges to understand the specific mean differences and statistical significance ($p < 0.05$).

All statistical analyses were regarded statistically significant when $p \leq 0.05$, and highly significant when $p < 0.001$.



4. RESULTS

4.1. Open Field Tests Results

We showed the open field test parameters as mean $\pm$ SE in Table 3.1.

Table 4. 1 Open Field Test Finding of Groups

	Group 1 ^a (n=12)	Group 2 ^b (n=12)	Group 3 ^c (n=12)	F	p
NEC	1.83 $\pm$ 0.71	1.25 $\pm$ 0.50 ^a	0.50 $\pm$ 0.67 ^a	1.65	0.20
STC	4.83 $\pm$ 1.23	1.50 $\pm$ 0.41 ^a	1.25 $\pm$ 0.50 ^a	6.12	0.005
STP	593.33 $\pm$ 2.42	598.50 $\pm$ 0.41 ^a	598.75 $\pm$ 0.50 ^a	4.44	0.02
NR	11.83 $\pm$ 1.08	18.33 $\pm$ 1.84 ^a	11.66 $\pm$ 1.56 ^b	6.14	0.005
ND	3.66 $\pm$ 0.41	2.50 $\pm$ 0.37	2.66 $\pm$ 0.37	2.61	0.08

Group 1=a: Sham group, Group 2=b: 400IU Vitamin D, Group 3=c: 1000IU Vitamin D, NEC: Number of entering to the central area, STC: Spent time in the central area, STP: Spent time in the peripheral area, NR: Number of rearing, ND: Number of defecations. (^a, this mean is different from Group 1)

There were significant differences among a group in term of open field test value except for NEC and ND (Table 4.1).




Figure 3. 4. Comparison of spent time in center and in peripheral between all groups.

STC values of Group 2 ($p < 0.02$) and Group 3 ($p < 0.01$) were statistically lower than group 1. On the contrary, the STP values of Group 2 ($p < 0.05$) and Group 3 ($p < 0.04$) were higher than group 1. While the STC value of Group 3 was the lowest, the STP value was nearly the same in Group 2 and Group 3 (Figure 3.4).

The NR value was the highest in Group 2. There was a difference between Group 1 and Group 2 ($p < 0.01$) and between Group 2 and group 3 ($p < 0.01$) in terms of NR.

There were no differences among the groups for NEC and ND.

4.2. Elevated Plus Maze Results

We showed the elevated plus-maze test findings as mean $\pm$ SE at Table 3.2.

Table 4. 2 The elevated plus-maze finding of groups (mean $\pm$ SE)

	Group 1 ^a (n=12)	Group 2 ^b (n=12)	Group 3 ^c (n=12)	F	p
STCA	217.16 $\pm$ 33.10	196.33 $\pm$ 19.98	234.58 $\pm$ 24.3	0.52	0.59
STOA	82.83 $\pm$ 33.1	103.66 $\pm$ 19.98	40.41 $\pm$ 12.22	1.89	0.16
ECA	1.50 $\pm$ 0.33	2.25 $\pm$ 0.37 ^c	1.16 $\pm$ 0.16	3.30	0.04
EOA	1.25 $\pm$ 0.21	1.33 $\pm$ 0.22	0.83 $\pm$ 0.24	1.38	0.26
Formula of Entry	47,46 $\pm$ 8,71	43.05 $\pm$ 6.09	29.16 $\pm$ 7.43	1.62	0.21
Formula of Time/Sec	25.00 $\pm$ 10.49	25.24 $\pm$ 5.85	13.47 $\pm$ 4.07	0.84	0.43

Group 1: Sham group, Group 2: 400 IU vitamin D, Group 3: 1000 IU vitamin D, STCA: Spent time in the closed arm, STOA: Spent time in open arm, ECA: Entry number of the closed arm, EOA: Entry number of open arms




Figure 3. 5. Comparison by percentage of spent time and number of enter in the open arms.

In Group 2, the STCA value was the lowest and the STOA value was the highest from other groups.

ECA value was statistically higher in Group 2 than Group 3. There were no statistical differences between groups for EOA value, for the formula of entry and for a formula of time among groups.



5. DISCUSSION

In our study, we wanted to determine the effect of Vitamin D on anxiety in rats by using a variety of doses. Because of the increasing amount of research that suggests a substantial link between VD insufficiency and emotional disorders such as sadness and anxiety, we thought that extra vitamin D usage would prevent anxiety. Where serum 25(OH) D deficiency might cause anxiety and depression symptoms, and lower serum 25(OH) D levels could result in severe anxiety at baseline in older people [73, 74] since vitamin D can be produced naturally by human skin under suitable conditions, many people would easily exceed their daily recommended dose by taking unnecessary supplements. This could lead to undesirable side effects and possibly anxiogenic effect as we will discuss later.

The investigation was conducted to observe how different doses of vitamin D could result in behavioural changes and anxiety in rats, through both open field and elevated plus-maze tests. The results will help us understand better how vitamin D could affect deeply our mood and psychological behaviors in the future.

By conducting the OFT, we could notice some behavioural changes among the three groups. Rats receiving both doses of vitamin D spent the most time in the peripheral area and preferred not to explore the central area and group 3 showed the least exploring activity compared with group 1 and group 2. The time spent in the peripheral area in OFT was used as a sign of anxiety [75]. These results are consistent with (Burne, Thomas HJ, et al. (2006), where unrestricted DVD-deficient rats spent more time on the wall sides in the open field and spent less time on its corners and centre than control rats [76]. Fedotova et al. (2017) assessed also the effectiveness of chronic administration of vitamin D using the different doses through the subcutaneous route. In his study, the administration was done once a day for 14 days consecutively. Contrary to our results, this test showed that only the doses of 1.0 and 2.5 mg/kg per day showed anxiolytic effects, keeping in mind that the rats were adults and had their ovaries removed [77]. While the rats that took higher-dose vitamin D (G 3) in the EPM test had the lowest number of entries in the open arm, rats prefer to stay in the enclosed arms of the maze and avoid the open arms. This is owing to rats' natural dislike to open spaces, as well as the maze's height [78]. These results are consistent with (Fu, Lin, et al., 2017) an animal study that discovered that adulthood vitamin D deficiency was linked to mood-changing behaviours, were found the time of entry into the open arms was reduced in VDD diet-fed mice than in control animals [79], Baranenko et al. (2019) demonstrated

that the treatment with VD3 (5.0 mg/kg) in ovariectomized rats significantly reduced anxiety-like behaviour in the EPM [80]. The dose difference between that study and ours may be due to the fact that the rats used were ovariectomized not normal rats. However, both vitamin D overdosing and deficiency can alter the social behaviour of the rats.

Vitamin D is important for the metabolism of both phosphate and calcium in living beings. The main purpose of vitamin D is to ensure that the body absorbs calcium. Because of its vital role in preserving the basic structure of the organism) i.e., musculoskeletal health, Previous research showed that vitamin D is one of the components that found its receptor across the body, even the brain in the same areas related to depression and anxiety [81], where they expressed that vitamin D, activates genes that regulate the immune system and releases neurotransmitters that influence brain functions [82]. Also, because vitamin D is the most important factor in calcium metabolism, high consumption of vitamin D can lead to hypercalcemia as it can increase the level of calcium in the blood [83].

Hypercalcemia caused by a high intake of vitamin D over a long period has been linked to neuropsychiatric problems. Patients with hypercalcemia usually have anxiety, but patients with severe hypercalcemia have changed mental status [84]. The mechanism of hypercalcemia-induced psychosis is still unknown. This can be explained as a result of variations in monoamine levels in the central nervous system and glutaminergic excitotoxicity via N-Methyl-D-aspartate receptors could explain it, though [85].

Vitamin D is generated in the brain and then passes through the blood-brain barrier to operating directly on cells that contain its nuclear receptor [86]. Vitamin D metabolites may alter the behaviour of cognitive functions through their particular neuroprotective effects, according to recent studies. In tests testing cognitive functioning, people with low vitamin D levels do worse [87], while others show that those with appropriate vitamin and micronutrient concentrations have improved cognitive skills [88].

According to some research, dopamine appears to have an essential role in anxiety modulation in several areas of the brain, according to studies. The mesolimbic, mesocortical, and nigrostriatal dopaminergic systems have been implicated in anxiety [89].

In certain studies, vitamin D has been found to have a neuroprotective impact on dopaminergic pathways in the adult brain. For example, when 6-hydroxydopamine (6-OHDA) is given to rats that have been pre-treated for one week with 1,25(OH) 2D3, the dopaminergic function in the treated rats is intact. [90]. Increased dopamine (DA) and

alterations in the caudate-putamen and hypothalamus were seen in the brainstem of rats given a single dose of vitamin D postnatal [91]. In vitro, 1,25(OH)₂D₃ promotes tyrosine hydroxylase production in adrenal medullary cells. [92] Vitamin D has been demonstrated to increase the production of tyrosine hydroxylase, hinting that it may have an impact on dopaminergic functions [93].

Serotonin 5-hydroxytryptamine or 5-HT is a neurotransmitter in 1948, it was discovered. that is hypothesized to play a role in pain sleep appetite anxiety depression and migraines among other things serotonin is made in the body from tryptophan which is converted to 5-HT by a variety of enzymes [94]. Serum serotonin levels increased significantly in the vitamin D-treated group compared to the non-treated group in a previous study this could be due to vitamin D's role in increased transcription of the tryptophan hydroxylase-2 gene which results in increased conversion tryptophan to serotonin in the brain [95]. As a result, in our study, we expect that high dose vitamin D intake might cause excess release of serotonin in the central nervous system and this release causes a disorder known as serotonin syndrome [96]. Serotonin poisoning is characterized by neuromuscular excitation altered mental state and autonomic stimulation [97].

6. CONCLUSION

According to our findings from open field test and elevated plus maze both different doses of Vitamin D showed anxiogenic effects. The mechanisms by which vitamin D increases anxiety are currently unknown, and additional research is required.

In the light of past findings and conclusions, this study's findings resulted in the following recommendations:

- 1- Use different vitamin D doses of low than 1000 IU are administered to experimental animals in order to determine the effects and changes that are expected to occur for anxiolytic effect.
- 2- To achieve better findings in future anxiolytic effect research, it is preferable to increase the number of experimental rats and shorten the experiment length to less than two months.

REFERENCES

1. Brown A, Dusso A. E. Slatopolsky. 1992 Vitamin D. The kidney: physiology and pathophysiology.:1505-51.
2. Mozaffari-Khosravi H, Nabizade L, Yassini-Ardakani SM, Hadinedoushan H, Barzegar K. The effect of 2 different single injections of high dose of vitamin D on improving the depression in depressed patients with vitamin D deficiency: a randomized clinical trial. *Journal of clinical psychopharmacology*. 2013 Jun 1;33(3):378-85.
3. Khundmiri SJ, Murray RD, Lederer E. PTH and vitamin D. *Comprehensive Physiology*. 2011 Jan 17;6(2):561-601.
4. Wilson LR, Tripkovic L, Hart KH, Lanham-New SA. Vitamin D deficiency as a public health issue: using vitamin D2 or vitamin D3 in future fortification strategies. *Proceedings of the Nutrition Society*. 2017 Aug;76(3):392-9.
5. DeLuca GC, Kimball SM, Kolasinski J, Ramagopalan SV, Ebers GC. The role of vitamin D in nervous system health and disease. *Neuropathology and applied neurobiology*. 2013 Aug;39(5):458-84.
6. Wrzosek M, Łukaszewicz J, Wrzosek M, Jakubczyk A, Matsumoto H, Piątkiewicz P, Radziwoń-Zaleska M, Wojnar M, Nowicka G. Vitamin D and the central nervous system. *Pharmacological reports*. 2013 Mar;65(2):271-8.
7. Sahota, O. (2014). Understanding vitamin D deficiency. *Age and ageing*, 43(5), 589.
8. Aquino SL, Cunha AT, Lima JG, Sena-Evangelista KC, Oliveira AG, Cobucci RN, Pedrosa LF. Effects of vitamin D supplementation on fasting glucose, dyslipidemia, blood pressure, and abdominal obesity among patients with metabolic syndrome: A protocol for systematic review and meta-analysis of randomized controlled trials. *Systematic Reviews*. 2020 Dec;9(1):1-6.
9. Blank S, Scanlon KS, Sinks TH, Lett S, Falk H. An outbreak of hypervitaminosis D associated with the overfortification of milk from a home-delivery dairy. *American Journal of Public Health*. 1995 May;85(5):656-9.
10. Schilling S. Epidemic vitamin D deficiency among patients in an elderly care rehabilitation facility. *Deutsches Ärzteblatt International*. 2012 Jan;109(3):33.

11. Bertone-Johnson ER, Powers SI, Spangler L, et al. Vitamin D intake from foods and supplements and depressive symptoms in a diverse population of older women. *Am J Clin Nutr.* 2011;94(4):1104-1112
12. Fallah M, Askari G, Asemi Z. Is vitamin D status associated with depression, Anxiety and Sleep Quality in Pregnancy: A Systematic Review. *Advanced Biomedical Research.* 2020;9.
13. Penckofer S, Byrn M, Adams W, Emanuele MA, Mumby P, Kouba J, Wallis DE. Vitamin D supplementation improves mood in women with type 2 diabetes. *Journal of diabetes research.* 2017 Sep 7;2017.
14. Vanchinathan V, Lim HW. A dermatologist's perspective on vitamin D. In *Mayo Clinic Proceedings* 2012 Apr 1 (Vol. 87, No. 4, pp. 372-380). Elsevier.
15. Ledesma-Amaro R, Santos MA, Jiménez A, Revuelta JL. Microbial production of vitamins. In *Microbial production of food ingredients, enzymes and nutraceuticals* 2013 Jan 1 (pp. 571-594). Woodhead Publishing.
16. Chen TC, Lu Z, Holick MF. Photobiology of vitamin D. In *Vitamin D* 2010 (pp. 35-60). Humana press.
17. HoRsT RL, Reinhardt TA, Reddy GS. Vitamin D metabolism. *Vitamin D.* 2005 Jan 1; 1:15-36.
18. Norman AW. From vitamin D to hormone D: fundamentals of the vitamin D endocrine system essential for good health. *The American journal of clinical nutrition.* 2008 Aug 1;88(2):491S-9S.
19. Jean G, Souberbielle JC, Chazot C. Vitamin D in chronic kidney disease and dialysis patients. *Nutrients.* 2017 Apr;9(4):328.
20. Bouillon R, Norman AW, Lips P. Vitamin D deficiency. *N Engl J Med.* 2007 Nov 8;357(19):1980-.
21. Lu Z, Chen TC, Zhang A, Persons KS, Kohn N, Berkowitz R, Martinello S, Holick MF. An evaluation of the vitamin D₃ content in fish: Is the vitamin D content adequate to satisfy the dietary requirement for vitamin D? *The Journal of steroid biochemistry and molecular biology.* 2007 Mar 1;103(3-5):642-4.
22. Whitton C, Nicholson SK, Roberts C, Prynne CJ, Pot GK, Olson A, Fitt E, Cole D, Teucher B, Bates B, Henderson H. National Diet and Nutrition Survey: UK food

- consumption and nutrient intakes from the first year of the rolling programme and comparisons with previous surveys. *British journal of nutrition*. 2011 Dec;106(12):1899-914.
23. Tylavsky FA, Cheng S, Lyytikainen A, Viljakainen H, Lamberg-Allardt C. Strategies to improve vitamin D status in northern European children: exploring the merits of vitamin D fortification and supplementation. *The Journal of nutrition*. 2006 Apr 1;136(4):1130-4.
 24. Ogan D, Pritchett K. Vitamin D and the athlete: risks, recommendations, and benefits. *Nutrients*. 2013 Jun;5(6):1856-68.
 25. Clarke HE, Coates ME, Eva JK, Ford DJ, Milner CK, O'donoghue PN, Scott PP, Ward RJ. Dietary standards for laboratory animals: report of the Laboratory Animals Centre Diets Advisory Committee. *Laboratory Animals*. 1977 Jan 1;11(1):1-28.
 26. Wacker M, Holick MF. Sunlight and Vitamin D: A global perspective for health. *Dermato-endocrinology*. 2013 Jan 1;5(1):51-108.
 27. Christakos S, Dhawan P, Verstuyf A, Verlinden L, Carmeliet G. Vitamin D: metabolism, molecular mechanism of action, and pleiotropic effects. *Physiological reviews*. 2016 Jan;96(1):365-408.
 28. Chun RF, Peercy BE, Orwoll ES, Nielson CM, Adams JS, Hewison M. Vitamin D and DBP: the free hormone hypothesis revisited. *The Journal of steroid biochemistry and molecular biology*. 2014 Oct 1; 144:132-7.
 29. Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *The Journal of nutrition*. 2005 Feb 1;135(2):317-22.
 30. Heaney RP. Assessing vitamin D status. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2011 Sep 1;14(5):440-4.
 31. Christakos S. In search of regulatory circuits that control the biological activity of vitamin D. *Journal of Biological Chemistry*. 2017 Oct 20;292(42):17559-60.
 32. Pekkinen M, Saarnio E, Viljakainen HT, Kokkonen E, Jakobsen J, Cashman K, Mäkitie O, Lamberg-Allardt C. Vitamin D binding protein genotype is associated with serum 25-hydroxyvitamin D and PTH concentrations, as well as bone health in children and adolescents in Finland. *PloS one*. 2014 Jan 30;9(1): e87292.

33. DeLuca HF. Overview of general physiologic features and functions of vitamin D. *The American journal of clinical nutrition*. 2004 Dec 1;80(6):1689S-96S.
34. Holick MF. McCollum Award Lecture, 1994: vitamin D—new horizons for the 21st century. *The American journal of clinical nutrition*. 1994 Oct 1;60(4):619-30.
35. Holick MF. Vitamin D deficiency. *New England journal of medicine*. 2007 Jul 19;357(3):266-81.
36. Wielders JP, Wijnberg FA. Preanalytical stability of 25 (OH)–vitamin D3 in human blood or serum at room temperature: Solid as a rock. *Clinical chemistry*. 2009 Aug 1;55(8):1584-5.
37. Cheng JB, Motola DL, Mangelsdorf DJ, Russell DW. De-orphanization of cytochrome P450 2R1: a microsomal vitamin D 25-hydroxylase. *Journal of Biological Chemistry*. 2003 Sep 26;278(39):38084-93.
38. De La Hunt A, Wallace AM, Gibson S, Viljakainen H, Lamberg-Allardt C, Ashwell M. UK Food Standards Agency Workshop Consensus Report: the choice of method for measuring 25-hydroxyvitamin D to estimate vitamin D status for the UK National Diet and Nutrition Survey. *British Journal of Nutrition*. 2010 Aug;104(4):612-9.
39. Zhu J, DeLuca HF. Vitamin D 25-hydroxylase—four decades of searching, are we there yet? *Archives of biochemistry and biophysics*. 2012 Jul 1;523(1):30-6.
40. Bouillon R. Comparative analysis of nutritional guidelines for vitamin D. *Nature Reviews Endocrinology*. 2017 Aug;13(8):466-79.
41. Hernández Rosiles V, Duarte Salazar C, Maldonado Velazquez R, Rivas Ruiz R, Clark P. Determination of 25 (OH) D serum levels in children with systemic lupus erythematosus and juvenile idiopathic arthritis. *Boletín Médico Del Hospital Infantil de México*. 2015 Apr;72(2):99-105.
42. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, Kovacs CS. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *The Journal of Clinical Endocrinology & Metabolism*. 2011 Jan 1;96(1):53-8.
43. Alshahrani F, Aljohani N. Vitamin D: deficiency, sufficiency and toxicity. *Nutrients*. 2013 Sep;5(9):3605-16.

44. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *The American journal of clinical nutrition*. 2004 Dec 1;80(6):1678S-88S.
45. Mawer EB, Hann JT, Berry JL, Davies M. Vitamin D metabolism in patients intoxicated with ergocalciferol. *Clinical Science*. 1985 Feb;68(2):135-41.
46. Cui X, Gooch H, Petty A, McGrath JJ, Eyles D. Vitamin D and the brain: Genomic and non-genomic actions. *Molecular and cellular endocrinology*. 2017 Sep 15; 453:131-43.
47. Isaac ML, Larson EB. Medical conditions with neuropsychiatric manifestations. *Medical Clinics*. 2014 Sep 1;98(5):1193-208.
48. Nagy L, Mangini P, Schroen C, Aziz R, Tobia A. Prolonged Hypercalcemia-Induced Psychosis. *Case reports in psychiatry*. 2020 Feb 1;2020.
49. Cherniack EP, Troen BR, Florez HJ, Roos BA, Levis S. Some new food for thought: the role of vitamin D in the mental health of older adults. *Curr Psychiatry Rep*. 2009 Feb 1;11(1):12-9.
50. Gezen-Ak D, Dursun E, Yilmazer S. The effect of vitamin D treatment on nerve growth factor (NGF) release from hippocampal neurons. *Nöro Psikiyatri Arşivi*. 2014 Jun;51(2):157.
51. Eyles DW, Smith S, Kinobe R, Hewison M, McGrath JJ. Distribution of the vitamin D receptor and 1 α -hydroxylase in human brain. *Journal of chemical neuroanatomy*. 2005 Jan 1;29(1):21-30.
52. Kazdin AE. *Encyclopedia of psychology*. American Psychological Association, editor. Washington, DC: American Psychological Association; 2000.
53. Lykouras L, Michopoulos J. Anxiety disorders and obesity. *Psychiatriki*. 2011 Oct 1;22(4):307-13.
54. Rodgers R, Cao BJ, Dalvi A, Holmes A. Animal models of anxiety: an ethological perspective. *Brazilian journal of medical and biological research*. 1997; 30:289-304.
55. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, Kovacs CS. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *The Journal of Clinical Endocrinology & Metabolism*. 2011 Jan 1;96(1):53-8.

56. Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *The American journal of clinical nutrition*. 2007 Jan 1;85(1):6-18.
57. Bays TB, Lightfoot T, Mayer J. *Exotic pet behavior: birds, reptiles, and small mammals*. Elsevier Health Sciences; 2006.
58. Asth L, Lobão-Soares B, André E, de Paula Soares V, Gavioli EC. The elevated T-maze task as an animal model to simultaneously investigate the effects of drugs on long-term memory and anxiety in mice. *Brain research bulletin*. 2012 Apr 10;87(6):526-33.
59. Ennaceur A, Chazot PL. Preclinical animal anxiety research—flaws and prejudices. *Pharmacology research & perspectives*. 2016 Apr;4(2): e00223.
60. Ari C, D’Agostino DP, Diamond DM, Kindy M, Park C, Kovács Z. Elevated plus maze test combined with video tracking software to investigate the anxiolytic effect of exogenous Ketogenic supplements. *JoVE (Journal of Visualized Experiments)*. 2019 Jan 7(143): e58396.
61. Fuchs E, Flügge G. Experimental animal models for the simulation of depression and anxiety. *Dialogues in clinical neuroscience*. 2006 Sep;8(3):323.
62. Uekita T, Ishibashi A, Sakamoto T. Influence of Pre-Weaning Social Isolation on Post-Weaning Emotion Tendency and Mother–Infant Interactions in Infant Octodon Degus. *International journal of environmental research and public health*. 2019 Jan;16(10):1824.
63. Kumar V, Bhat ZA, Kumar D. Animal models of anxiety: a comprehensive review. *Journal of pharmacological and toxicological methods*. 2013 Sep 1;68(2):175-83.
64. Graeff FG, Viana M, Tomaz C. The elevated T maze, a new experimental model of anxiety and memory: effect of diazepam. *Brazilian journal of medical and biological research= Revista brasileira de pesquisas medicas e biologicas*. 1993 Jan 1;26(1):67-70.
65. Seibenhener ML, Wooten MC. Use of the open field maze to measure locomotor and anxiety-like behavior in mice. *JoVE (Journal of Visualized Experiments)*. 2015 Feb 6(96):e52434.
66. Kraeuter AK, Guest PC, Sarnyai Z. The elevated plus maze test for measuring anxiety-like behavior in rodents. In *Pre-clinical models 2019* (pp. 69-74). Humana Press, New York, NY.

67. Weiss JM, Cierpial MA, West CH. Selective breeding of rats for high and low motor activity in a swim test: toward a new animal model of depression. *Pharmacology Biochemistry and Behavior*. 1998 Sep 1;61(1):49-66.
68. Gellért L, Varga D. Locomotion activity measurement in an open field for mice. *Bio-protocol*. 2016 Jul 5;6(13): e1857-.
69. Šimoliūnas E, Rinkūnaitė I, Bukelskienė Ž, Bukelskienė V. Bioavailability of different vitamin D Oral supplements in laboratory animal model. *Medicina*. 2019 Jun;55(6):265.
70. Rooney MR, Harnack L, Michos ED, Ogilvie RP, Sempos CT, Lutsey PL. Trends in use of high-dose vitamin D supplements exceeding 1000 or 4000 international units daily, 1999-2014. *Jama*. 2017 Jun 20;317(23):2448-50.
71. Braun AA, Skelton MR, Vorhees CV, Williams MT. Comparison of the elevated plus and elevated zero mazes in treated and untreated male Sprague–Dawley rats: effects of anxiolytic and anxiogenic agents. *Pharmacology Biochemistry and Behavior*. 2011 Jan 1;97(3):406-15.
72. Suchecki D, Tiba PA, Tufik S. Hormonal and behavioural responses of paradoxical sleep-deprived rats to the elevated plus maze. *Journal of neuroendocrinology*. 2002 Jul;14(7):549-54.
73. Zhu, C., Zhang, Y., Wang, T., Lin, Y., Yu, J., Xia, Q., ... Zhu, D. (2020). Vitamin D supplementation improves anxiety but not depression symptoms in patients with vitamin D deficiency. *Brain and Behavior*, 10(11),
74. Bicikova M, Duskova M, Vitku J, Kalvachová B, Ripova D, Mohr P, Stárka L. Vitamin D in anxiety and affective disorders. *Physiological research*. 2015;64: S101.
75. Schmitt U, Hiemke C. Strain differences in open-field and elevated plus-maze behavior of rats without and with pretest handling. *Pharmacology Biochemistry and Behavior*. 1998 Apr 1;59(4):807-11.
76. Burne TH, O'Loan J, McGrath JJ, Eyles DW. Hyperlocomotion associated with transient prenatal vitamin D deficiency is ameliorated by acute restraint. *Behavioural brain research*. 2006 Nov 1;174(1):119-24.
77. Fedotova J, Pivina S, Sushko A. Effects of chronic vitamin D3 hormone administration on anxiety-like behavior in adult female rats after long-term ovariectomy. *Nutrients*. 2017 Jan;9(1):28.

78. Tarbali S, Khezri S. Protective Effects of Vitamin D3 on Anxiety-like Behavior and the Total Antioxidant Power Following the Local Injection of Lysophosphatidylcholine in the Adult Rat Dorsal Hippocampus. *Journal of Neurological Sciences*. 2015 Jul 1;32(3).
79. Fu L, Chen YH, Chen X, Xu S, Yu Z, Xu DX. Vitamin D deficiency impairs neurobehavioral development in male mice. *Physiology & behavior*. 2017 Oct 1; 179:333-9.
80. Baranenko D, Fedotova J, Van Den Tol AJ. Vitamin D 3 attenuates anxiety-like behavior in long-term ovariectomized rats with unpredictable mild stress. *Biocell*. 2019;43(4):299.
81. Roy NM, Al-Harhi L, Sampat N, Al-Mujaini R, Mahadevan S, Al Adawi S, Essa MM, Al Subhi L, Al-Balushi B, Qoronfleh MW. Impact of vitamin D on neurocognitive function in dementia, depression, schizophrenia and ADHD. *Front Biosci (Landmark Ed)*. 2021 Jan 1; 26:566-611.
82. Greenblatt JM. Psychological consequences of vitamin D deficiency. *Psychology today*. 2011 Nov;62.
83. Guerra V, Vieira OM, Laurindo AF, Paula FJ, Moysés M. Hypercalcemia and renal function impairment associated with vitamin D toxicity: case report. *Brazilian Journal of Nephrology*. 2016 Oct; 38:466-9.
84. Isaac ML, Larson EB. Medical conditions with neuropsychiatric manifestations. *Medical Clinics*. 2014 Sep 1;98(5):1193-208.
85. Nagy L, Mangini P, Schroen C, Aziz R, Tobia A. Prolonged Hypercalcemia-Induced Psychosis. *Case reports in psychiatry*. 2020 Feb 1;2020.
86. Kozyra M, Zimnicki P, Kaczerska J, Śmiech N, Nowińska M, Milanowska J. The effect of vitamin D on mental health—literature analysis. *Journal of Education, Health and Sport*. 2020 Aug 29;10(8):398-407.
87. Przybelski RJ, Binkley NC. Is vitamin D important for preserving cognition? A positive correlation of serum 25-hydroxyvitamin D concentration with cognitive function. *Archives of Biochemistry and Biophysics*. 2007 Apr 15;460(2):202-5.
88. Bryan J, Osendarp S, Hughes D, Calvaresi E, Baghurst K, van Klinken JW. Nutrients for cognitive development in school-aged children. *Nutrition reviews*. 2004 Aug 1;62(8):295-306.

89. Siega-Riz AM, Popkin BM. Dietary trends among low socioeconomic status women of childbearing age in the United States from 1977 to 1996: a comparison among ethnic groups. *JOURNAL-AMERICAN MEDICAL WOMENS ASSOCIATION*. 2001 Jan 1;56(2):44-8.
90. Zarrindast MR, Khakpai F. The modulatory role of dopamine in anxiety-like behavior. *Archives of Iranian medicine*. 2015 Sep 1;18(9):0-.
91. Eyles DW, Feron F, Cui X, Kesby JP, Harms LH, Ko P, McGrath JJ, Burne TH. Developmental vitamin D deficiency causes abnormal brain development. *Psychoneuroendocrinology*. 2009 Dec 1;34: S247-57.
92. Tekes K, Gyenge M, Hantos M, Csaba G. Transgenerational hormonal imprinting caused by vitamin A and vitamin D treatment of newborn rats. Alterations in the biogenic amine contents of the adult brain. *Brain and Development*. 2009 Oct 1;31(9):666-70.
93. Kesby JP, Eyles DW, Burne TH, McGrath JJ. The effects of vitamin D on brain development and adult brain function. *Molecular and cellular endocrinology*. 2011 Dec 5;347(1-2):121-7.
94. Puchacz E, Stumpf WE, Stachowiak EK, Stachowiak MK. Vitamin D increases expression of the tyrosine hydroxylase gene in adrenal medullary cells. *Molecular Brain Research*. 1996 Feb 1;36(1):193-6.
95. Dunkley EJ, Isbister GK, Sibbritt D, Dawson AH, Whyte IM. The Hunter Serotonin Toxicity Criteria: simple and accurate diagnostic decision rules for serotonin toxicity. *Qjm*. 2003 Sep 1;96(9):635-42.
96. Isbister GK, Buckley NA, Whyte IM. Serotonin toxicity: a practical approach to diagnosis and treatment. *Medical Journal of Australia*. 2007 Sep;187(6):361-5.
97. Volpi-Abadie J, Kaye AM, Kaye AD. Serotonin syndrome. *Ochsner Journal*. 2013 Dec 21;13(4):533-40.

APPENDIX (I)



1993

BAŞKENT ÜNİVERSİTESİ
Tıp ve Sağlık Bilimleri Araştırma Kurulu



Sayı : 94603339-604.01.02/ 33348
Konu : Proje Onayı

18/11/2020

FİZYOLOJİ ANABİLİM DALINA

Fizyoloji Anabilim Dalında görev yapmakta olan Prof. Dr. Nazan Dolu'nun danışmanlığında Sağlık Bilimleri Enstitüsü / Fizyoloji Tezli Yüksek Lisans Programı öğrencisi Zahour Gamal Eddn Habib Asmaeil'in sorumluluğunda yürütülecek olan DA20/19 nolu "Effects of high doses of vitamin D on anxiety and exploratory activity behaviours in rats" başlıklı araştırma projesi Kurulumuz ve Hayvan Deneyleri Yerel Etik Kurulunun 02/11/2020 tarih ve 20/17 sayılı kararı ile uygun görülmüştür. Projenin başlama tarihi ile çalışmanın sunulduğu kongre ve yayımlandığı dergi konusunda Kurulumuza bilgi verilmesini rica ederim.

e-imzalıdır

Kurul Başkanı

Not: Çalışma bildiri ve/veya makale haline geldiğinde "Gereç ve Yöntem" bölümüne aşağıdaki ifadelerden uygun olanının eklenmesi gerekmektedir.

— Bu çalışma Başkent Üniversitesi Hayvan Deneyleri Yerel Etik Kurulu tarafından onaylanmış (Proje no:...) ve Başkent Üniversitesi Araştırma Fonunca desteklenmiştir.

— This study was approved by Baskent University Ethical Committee for Experimental Resarch on Animals (Project no:...) and supported by Baskent University Research Fund.

DAĞITIM

Sağlık Bilimleri Enstitüsü Müdürlüğüne
Fizyoloji Anabilim Dalına

APPENDIX (2)



1993
BAŞKENT UNIVERSITY

LOCAL ETHICS COMMITTEE FOR ANIMAL EXPERIMENTS DECISION		
SESSION NO	DECISION NO	DATE OF DECISION
09	20/17	02/11/2020

Project DA20/19 no entitled “Effects of high doses of vitamin D on anxiety and exploratory activity behaviours in rats” pending to be conducted by Nazan Dolu with the Department of Physiology has been reviewed and unanimously approved by the Local Ethics Committee for Animal Experiments.