



REPUBLIC OF TÜRKİYE
ALTINBAŞ UNIVERSITY
Institute of Graduate Studies
Biomedical Sciences

**ANTIMICROBIAL AND ANTI-BIOFILM
ACTIVITIES OF LIPIDS EXTRACTED FROM
SCHIZOCHYTRIUM SP. S31 MICROALGAE**

Doaa Abdullah Hammadi AL-OGAIDI

Master's Thesis

Supervisor

Asst. Prof. Dr. Nurcan Vardar Yel

Istanbul, 2022

**ANTIMICROBIAL AND ANTI-BIOFILM ACTIVITIES OF LIPIDS
EXTRACTED FROM *SCHIZOCHYTRIUM* SP. S31 MICROALGAE**

Doaa Abdullah Hammadi AL-OGAIDI

Biomedical Sciences

Master's Thesis

ALTINBAŞ UNIVERSITY

2022

The thesis/dissertation titled ANTIMICROBIAL AND ANTI-BIOFILM ACTIVITIES OF LIPIDS EXTRACTED FROM *SCHIZOCHYTRIUM* SP. S31 MICROALGAE prepared by DOAA ABDULLAH HAMMADI AL-OGAIDI and submitted on 20/12/2022 has been **accepted unanimously** for the degree of Master of Science in Biomedical Science.

Assoc. Prof. Dr. Rafig GURBANOV

Co-Supervisor

Assist. Prof. Dr. Nurcan Vardar Yel

Supervisor

Thesis Defense Committee Members:

Assist. Prof. Dr. Zeliha Doğan

Vocational School of Health
Services, First and Emergency
Aid Program, Altınbaş
University

Assist. Prof. Dr. Ahmet Murat Şenışık

Vocational School of Health
Services, Radiotherapy
Program, Altınbaş University

Assist. Prof. Dr. Havva Esra Tütüncü

Faculty of Health Sciences,
Nutrition and Dietetics,
Malatya Turgut Özal
University

I hereby declare that this thesis/dissertation meets all format and submission requirements of a Master's thesis/dissertation.

Submission date of the thesis to the Graduate Education Institute: ____/____/____

I hereby declare that all information/data presented in this graduation project has been obtained in full accordance with academic rules and ethical conduct. I also declare all unoriginal materials and conclusions have been cited in the text and all references mentioned in the reference list have been cited in the text, and vice versa as required by the abovementioned rules and conduct.

Doaa Abdullah Hammadi AL-OGAIDI

Signature

DEDICATION

To my parents, for their endless love, support, and encouragement.

To my grandfather Hamed

To my brothers Taha and Yaseen.

To my sisters Hind, Huda, Sabaa, and Ghofran.

To my teachers

Doaa Abdullah Hammadi AL-OGAIDI

ACKNOWLEDGEMENTS

I thank God for giving me this gift and giving me the strength to reach this point and complete it in the best way.

I thank the department and its staff, headed by Prof. Dr. Feride Severcan for the effort in facilitating my task to complete it in the best way.

I would like to express my sincere gratitude to my supervisors Assist. Prof. Dr. Nurcan VARDAR YEL and Assoc. Prof. Dr. Rafiq GURBANOV for their continuous support, motivation, and immense knowledge.

I do not forget to thank my close friends who spared no effort to help me. In addition to everyone who helped me, even if it was a simple thing.

Last but not least, my big family at a time full of love and cooperation, the International Academy for Leadership and Development (IALD) and a big thanks to Mrs. Mona AL-KATEEB and Mr. Abdulrahman AL-AHMAD.

ABSTRACT

ANTIMICROBIAL AND ANTI-BIOFILM ACTIVITIES OF LIPIDS EXTRACTED FROM *SCHIZOCHYTRIUM* SP. S31 MICROALGAE

AL-OGAIDI, Doaa Abdullah Hammadi

M.Sc. Biomedical Science, Altınbaş University, Turkey

Supervisor: Assist. Prof. Dr. Nurcan VARDAR YEL

Co-Supervisor: Assoc. Prof. Dr. Rafiq GURBANOV

Date: December/2022

Pages: 92

Biotechnology has been developing tools and procedures for the benefit of living things for many years. In most of these studies, the production of essential metabolic compounds is prioritized for human health. One of the crucial substances for human health that has gained popularity recently is omega-3 fatty acids. Alternative sources of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have been searched since fish oils, the main commercial sources of omega-3 fatty acids, have heavy metal content, especially mercury, limited availability, and an unpleasant odor. *Schizochytrium* sp. S31 is a microalga with a high concentration of polyunsaturated fatty acids, particularly DHA. Because of its high DHA content and low heavy metal levels, heterotrophic cultivation of *Schizochytrium* sp. S31 provides an excellent opportunity to provide alternative omega-3 sources. It also contains a variety of bioactive substances that can be utilized as medicinal raw materials, food additives, animal feed, biodiesel and antimicrobial agents.

In this study, optimum conditions were applied to cultivate *Schizochytrium* sp. S31 heterotrophically and the nutritional composition of the dry biomass were analysed. According to the results, 45% protein, 5% total fat and 37% carbohydrate were obtained. Downstream processes, including cell lysis and lipid extraction, were adjusted in order to increase overall lipid content of *Schizochytrium* sp. S31. Five distinct solvent mixtures—hexane, hexane/ethanol (2:1), ethanol, methanol, and hexane/methanol (2:1)—as well as two different cell lysis techniques—mortar and pestle and ultrasonication—were examined and compared. Based on the results of cell lysis and lipid extraction, ultrasonication of hexane /methanol (2:1) enhanced the total lipid yields to 34% with a transparent appearance. According to the GCMS analysis results, 10.4% DHA and 16.11% EPA fatty acids were obtained. Total Omega-3 Fatty Acids and Total Omega-3 + Omega-6 Fatty Acids ratio were calculated as 27.42% and 32.15% respectively.

The antibacterial effects of *Schizochytrium* sp. S31 oil samples were also investigated on Gram-negative pathogens, *Pseudomonas aeruginosa*, *Escherichia coli* 35218 and *Klebsiella pneumonia*, which cause many different human infections, as well as the Gram-positive pathogen *Streptococcus mutans*. At the same time, their effects on biofilm formation in *Pseudomonas aeruginosa*, *Escherichia coli* 35218 bacteria were investigated. The *Schizochytrium* sp. S31 oil sample was found to have anti-pathogenic activity on pathogenic bacteria of *Pseudomonas aeruginosa*, *Escherichia coli* 35218, *Klebsiella pneumonia* and *Streptococcus mutans*. The surface adhesion ability and biofilm formation capability of *P. aeruginosa* and *Escherichia coli* 35218 strain, which are widely employed as model bacteria in biofilm research, were dramatically inhibited in the presence of a *Schizochytrium* sp. S31 oil sample.

Keywords: *Schizochytrium* sp. S31, antibacterial activity, antibiofilm, fatty acid, docosahexaenoic acid.

TABLE OF CONTENTS

	<u>Pages</u>
ABSTRACT.....	vii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
ABBREVIATIONS	Hata! Yer işareti tanımlanmamış.
1. INTRODUCTION AND PURPOSE.....	Hata! Yer işareti tanımlanmamış.
2. GENERAL INFORMATION.....	3
2.1 CLASSIFICATION OF ALGAE	3
2.1.1 Macroalgae.....	3
2.1.2 Cyanobacteria.....	4
2.1.3 Microalgae	5
2.2 IMPORTANT MICROALGAE SPECIES	6
2.2.1 <i>Spirulina</i>	6
2.2.2 <i>Chlorella</i>	7
2.2.3 <i>Dunaliella</i>	7
2.2.4 <i>Haematococcus</i>	8
2.2.5 <i>Schizochytrium</i> sp.....	9
2.3 COMPOSITION OF MICROALGAL BIOMASS.....	10
2.3.1 Carbohydrates	11

2.3.2 Protein	12
2.3.3 Lipids	12
2.4 LONG CHAIN POLYUNSATURATED FATTY ACIDS (PUFA).....	13
2.4.1 Omega-3	14
2.4.1.1 Eicosapentaenoic acid(EPA)	15
2.4.1.2 Docosaheaxaenoic acid(DHA)	16
2.5 ANTIMICROBIAL ACTIVITY OF MICROALGAE.....	16
2.5.1 Antibacterial Activity	18
2.5.2 Antiviral Activity.....	21
2.5.3 Antifungal Activity	22
2.6 ALGAL FATTY ACIDS AS ANTIMICROBIAL COMPOUND.....	24
2.7 ANTIBIOFILM ACTIVITY	27
3. METHODOLOGY	30
3.1 MICROORGANISM AND MEDIUMS	30
3.2 GROWTH EXPERIMENT	30
3.3 DOWNSTREAM PROCESS.....	30
3.3.1 Harvesting The Microalgal Culture	30
3.3.2 Lyophilization	31
3.3.3 Cell Disruption	31
3.3.3.1 Mortar and pestle	31
3.3.3.2 Sonication.....	32

3.4 DRY BIOMASS ANALYSIS.....	32
3.5 LIPID EXTRACTION.....	32
3.6 LIPID CHARACTERIZATION.....	32
3.6.1 Total Lipid Analysis	32
3.6.2 Fatty Acid Analysis	33
3.6.3 Gas Chromatography Analysis	33
3.7 ANTIBACTERIAL ACTIVITY ASSAY.....	33
3.8 BIOFILM FORMATION ASSAY	34
4. RESULTS	35
4.1 GROWTH EXPERMENTS.....	35
4.2 DRY BIOMASS CALCULATION AND ANALYSIS	36
4.3 COMPARISON OF SEVERAL CELL LYSIS AND LIPID EXTRACTION TECHNIQUES	38
4.4 GAS CHROMATOGRAPHY ANALYSIS	39
4.5 ANTIBACTERIAL ACTIVITY.....	41
4.6 ANTIBIOFILM ACTIVITY.....	46
5. DISCUSSION AND CONCLUSION	50
REFERENCES.....	54
APPENDIX A.....	71

LIST OF TABLES

	<u>Pages</u>
Table 2.1: Antibacterial properties of compounds from microalgae	19
Table 2.2: Antiviral properties of compounds from microalgae.....	21
Table 2.3: Antifungal properties of compounds from microalgae.....	23
Table 2.4: A list of bioactive molecules isolated from different algal species.....	24
Table 2.5: Selected compounds from various microalgae and their antibiofilm.....	28
Table 4.1: Chemical characteristics of the of <i>Schizochytrium</i> sp.S31 dry biomass.....	36
Table 4.2: Total amino acid content of the dry biomass	37
Table 4.3: Total vitamin content of the dry biomass	38
Table 4.4: Fatty acid composition of <i>Schizochytrium</i> sp. S31.....	40

LIST OF FIGURES

	<u>Pages</u>
Figure 2.1: Images of different cyanobacteria species.....	4
Figure 2.2: Microalgae under the microscope unicellular green algae with large cells	5
Figure 2.3: Spirulina meneghiniana algae under microscope	6
Figure 2.4: Chlorella under microscope.....	7
Figure 2.5: Dunaliella under microscope.....	8
Figure 2.6: <i>Haematococcus</i> sp. under microscope	9
Figure 2.7: <i>Schizochytrium</i> sp. Cells under microscope	10
Figure 2.8: Molecular structures of EPA and DHA fatty acids	16
Figure 4.1: Growth curve of <i>Schizochytrium</i> sp. S31	35
Figure 4.2: Comparative results of different cell lysis and lipid extraction methods	39
Figure 4.3: Investigation of antibacterial effect of <i>Schizochytrium</i> sp. S31 oil sample on <i>P. aeruginosa</i> bacteria	42
Figure 4.4: Investigation of antibacterial effect of <i>Schizochytrium</i> sp. S31 oil sample on <i>Escherichia coli</i> 35218.....	43
Figure 4.5: Investigation of antibacterial effect of <i>Schizochytrium</i> sp. S31 oil sample on <i>Streptococcus mutans</i>	44
Figure 1.6: Investigation of antibacterial effect of <i>Schizochytrium</i> sp. S31 oil sample on <i>Klebsiella pneumoniae</i>	45
Figure 4.7: Effect of <i>Schizochytrium</i> sp. S31 oil sample on biofilm formation in <i>P. aeruginosa</i>	47

Figure 4.8: Effect of <i>Schizochytrium</i> sp. S31 oil sample on biofilm formation in <i>Escherichia coli</i>	
.....	48



ABBREVIATIONS

ALA	: Alpha linolenic acid
ARA	: Arachidonic acid
ATCC	: American Type Culture Collection
CDW	: Cell Dry Weight
DHA	: Docosahexaenoic Acid
DMSO	: Dimethyl Sulfoxide
EPA	: Eicosapentaenoic Acid
FAME	: Fatty Acid Methyl Ester
FAO	: Food and Agriculture Organization
FFA	: Free Fatty Acid
GC	: Gas Chromatography
GCMS	: Gas Chromatography Mass Spectrometry
GLA	: Gamma linolenic acid
HPLC	: High Pressure Liquid Chromatography
MSG	: Monosodium Glutamate
PUFA	: Polyunsaturated Fatty Acids
TAG	: Triacylglycerol
TFA	: Total Fatty Acid
WHO	: World Health Organization

1. INTRODUCTION AND PURPOSE

Biotechnological and multidisciplinary research are now underway in quest of new advances that may be used for the benefit of all living things. The basic metabolic requirements for our metabolism are one of the most discussed topics globally since the human being takes precedence over all other considerations and health issues are given top importance. For instance, the percentage of DHA produced for supplemental feeding has dramatically increased during the early 2000s (Harun et al., 2010).

Algae are significant sources of a broad range of complex lipids that may be used in culinary, cosmetic, and pharmaceutical sectors (Hossain et al., 2005; Abd El-Baky & El-Baroty, 2013). It has been investigated whether microalgae can be utilized to create food supplements, lipids, enzymes, polymers, among other things. Some of these have been accomplished by the cultivation of microalgae in a variety of media and substrates (Perez-Garcia et al., 2011). Several studies have shown that marine macroalgae lipids have antibacterial, antiviral and cancer-fighting properties (Gerasimenko et al., 2010).

Long-chain polyunsaturated fatty acids (LC-PUFAs) are essential for human nutrition and brain development. Among all LC-PUFAs, DHA and EPA (Eicosapentaenoic acid) are vital for human growth. There is evidence that DHA helps the evolution of the human brain by regulating metabolisms such as sperm retina and brain lipids (Apt & Behrens, 1999; Chanda et al., 2018). Omega-3 fatty acids are essential for the immune system well as cardiovascular and neurological system issues (Peltomaa et al., 2018; Simopoulos, 2002). In addition, dietary DHA consumption affects amyloid precursor protein processing and reduces the risk of Alzheimer's disease by inhibiting amyloid formation (Lim et al., 2005). Furthermore, fatty acids have been investigated as possible therapeutic antimicrobial agents due to their potency, wide range of activity, and absence of typical resistance mechanisms (Desbois & Smith, 2010; Thormar, 2010; Mullen et al., 2010; P. Desbois, 2012; Feldlaufer et al., 1993; Zhang et al., 2012)

The aim of this study was to analyze the antibacterial and antibiofilm effect of lipid extracted from DHA-rich *Schizochytrium* sp. S31 microalgae. The antibacterial effects of *Schizochytrium* sp. S31 oil samples were studied on Gram-negative pathogens, *Pseudomonas aeruginosa*, *Escherichia coli* 35218 and *Klebsiella pneumonia*, which cause many different human infections, as well as the

Gram-positive pathogen *Streptococcus mutans*. At the same time, antibiofilm activity of the *Schizochytrium* sp. S31 oil on *Pseudomonas aeruginosa*, and *Escherichia coli* 35218 bacteria were investigated.



2. GENERAL INFORMATION

2.1 CLASSIFICATION OF ALGAE

Classification is the systematic grouping of organisms into groups that are based on evolutionary or structural links. The classification of algae into taxonomic groups may vary depending on many different factors. It is not possible to classify algae in terms of a single feature. Some algae live in a terrestrial environment, while others live in an aquatic environment. There are unicellular algae as well as multicellular algae. Algae can also be divided into prokaryotic and eukaryotic species. Nowadays, algae are commonly classified into three groups: macroalgae, cyanobacteria and microalgae (Baweja & Sahoo, 2015).

2.1.1 Macroalgae

Macroalgae (seaweed) are photosynthetic organisms that exist in aquatic habitats all over the globe, including the Arctic zone. Seaweeds have an important ecological and economic purpose. The highest variety is found on rocky seashores and coral reefs. Algae are thought to capture ten times the amount of solar energy as terrestrial plants. In food chains, they are the principal primary producers of organic molecules (Kaur et al., 2002).

Algae may be aquatic or subaerial, depending on whether they are on the surface or at the bottom of the water. Aquatic algae can tolerate high pH, temperature, turbidity, oxygen, and carbon dioxide concentrations. Subaerial (aerobic) algae are those that live in the air, soil, and litter, or water surface. There are several different types of algae, including epiphytic (found on living plants), epiphyllous (found on leaf surfaces), corticolous (found on bark, trunks, or stems), epizoic (found on animals), as well as lithophile (found in stone, brick, or cement), epixylous (found on dead wood, such as poles and posts), and epipetalous (found on metal surfaces). The growth habits of marine algae may be classified as supralittoral (above high tide), intertidal (onshores exposed to tidal cycles), or sublittoral (benthic environment from extreme low water to about 200 m deep in very clear water). The word "seaweed" typically refers to multicellular red, green, and brown marine algae. Each group is different in colour due to having different photosynthetic pigments. Red algae are known as Rhodophyta; brown algae are known as Ochrophyta; and green algae are

known as Chlorophyta. Marine plants, like terrestrial plants, are benign and should be preserved for future generations (Kaur et al., 2002).

2.1.2 Cyanobacteria

Cyanobacteria are naturally occurring photosynthetic bacteria found in all environments on Earth. However, although they may sometimes form colonies, they are normally unicellular (Figure 2.1). The oxygenation of the early earth's atmosphere and seas during the massive oxidation event 2.4 billion years ago is assumed to have been caused by cyanobacteria. They were previously known as Cyanophyta or blue-green algae because they are photosynthetic organism (Damatac & Cao, 2022). Iron is a crucial metal in the cosmos, and it plays a vital role in the development of cyanobacteria. The magnetic field created by the liquid iron in the earth's core allows life to exist on our planet. Ferredoxins (iron-sulfur proteins) are believed to have contributed to the development of fermentative bacteria (Rath et al., 2015). Certain archaea and bacteria share a common ancestor that can convert Fe (III) to Fe (II), which conserve energy and promotes growth. It is also thought that the ancestors of Proteobacteria and Cyanobacteria utilised Fe (OH⁺) as a significant electron donor for CO₂ fixation (Rath et al., 2015).




Figure 2.1: Images of different cyanobacteria species (Phiri, 2021).

The endosymbiotic theory states that endosymbiosis is the mechanism by which chloroplasts in eukaryotic plants and algae developed from cyanobacterial progenitors. Thermophilic Cyanobacteria can grow at over 45°C and were less common than mesophilic Cyanobacteria (Whitton, 2012). The maximum temperature limit for Cyanobacteria species in hot water is approximately 73°C, and *Synechococcus* sp. were discovered. (Damatac & Cao, 2022; Strunecký et al., 2019).

2.1.3 Microalgae

Microalgae are microscopic organisms that exist singly or in groups in the water and sediment (Figure 2.2). Similar to terrestrial plants, microalgae utilise the oxygen produced as a by-product of photosynthesis to produce organic molecules.




Figure 2.2: Microalgae under the microscope, unicellular green algae with large cells (Maltsev,2022).

Algae are utilized as food and fuel, and their extracts have significance in the medicinal business. These primitive plants have chlorophyll but no roots or leaves. Photosynthetic microalgae can use sunlight for energy to convert CO₂ into biomass and produce oxygen. Although microalgae develop in several ways, they can often double their biomass in 24 hours. There are around 50.000 microalgae species; however, only 30.000 have been examined (Handayania & Ariyantib, 2011).

2.2 IMPORTANT MICROALGAE SPECIES

The biotechnological use of microalgae has been concentrated on important species during the past two decades. Some of the important microalgae species are *Spirulina*, *Chlorella vulgaris*, *Dunaliella. salina*, *Haematococcus pluvialis*, and *Schizochytrium sp.* (McCann et al., 2014). The following subsections discuss the features, manufacturing method, and makeup of each of these microalgal species.

2.2.1 Spirulina

Spirulina is a symbiotic, filamentous, and multicellular blue-green microalgae that utilizes nitrogen from the surrounding environment. It is available in two separate shapes: spiral rod as shown in (Figure 2.3) or disk-shaped. The blue pigment phycocyanin serves as the principal photosynthetic pigment in Spirulina. Some include phycoerythrin, a pigment that gives microalgae their characteristic red or pink colour.




Figure 2.3: *Spirulina meneghiniana* algae under the microscope (Fox, 2022).

Spirulina is a well-studied algae for outdoor cultivation on a huge scale. It thrives in situations with high bicarbonate concentrations and pH levels (Paniagua-Michel et al., 2015; De Oliveira et al., 1999). Spirulina is cultured in raceway ponds powered by paddle wheels. The pond's water depth typically ranges between 300-500 mm, depending on the microalgal density and season. The water depth of a pond is also depended on its size as well as flow rate, and the algal culture's optimum light absorption. The temperature has a significant influence on spirulina production. It thrives at temperatures between 35°C and 37°C (De Oliveira et al., 1999).

2.2.2 Chlorella

Chlorella microalgae are photoautotrophic, single-celled, spherical (diameter: 5–10µm) and green in colour with no flagella (Figure 2.4) (Becker, 2007). It needs CO₂, water, light, and a few minerals in order to grow. Photobioreactors, large round tanks, mixed open ponds with paddle wheels, or circular open ponds have all been used commercially to cultivate Chlorella (Borowitzka & Larkum, 1987; Mobin et al., 2001). First, it is cultured in tiny containers and then transferred to a larger tank or pond.




Figure 2.4: *Chlorella* under microscope (Thongdumhyu, 2018).

Circular tanks and ponds are two of the most frequent outdoor cultivation techniques. The biggest pond has a surface area of (500 m²) and a depth of (200 mm), respectively. Chlorella microalgae for aquaculture are often produced on a smaller scale, such as in 20–40 L carboys or enormous plastic bags. (Borowitzka & Larkum, 1987). Centrifugation and autoflocculation are used to extract chlorella. Several ways to process the biomass after harvesting, including spray drying or drum drying. In terms of dry weight, Chlorella includes 2-46 % of lipid, 12-28 % of carbohydrates, and 11–58 % of proteins (Zhu et al., 2014). Chlorella is rich in vitamins, including provitamin A, carotene, retinol, vitamin E, and vitamin C (Borowitzka, 1998).

2.2.3 Dunaliella

They are green microalgae that are edible, nutrient-dense, single-cell, and flagellated as shown under the microscope in Figure2.5. Additionally, it is found in a variety of marine and freshwater settings. Dunaliella salina has garnered considerable interest owing to its remarkable antioxidant activity. It is one the best-known sources of high-carotenoid-containing compounds compared to

other sources, as it has β -carotene concentrations of up to 14% of dry biomass. Although *Dunaliella* can grow at salinities above 30% NaCl saturation, the ideal growing salinity is 22% NaCl saturation. *D. salina* is often farmed in huge (5–200 ha) shallow, shallow paddlewheel raceway ponds or unstirred ponds with an average size of less than 1000 m². This species thrives on a semi-continuous cultivation procedure (Borowitzka & Vonshak, 2017). Borowitzka et al. (Borowitzka & Vonshak, 2017) showed that *Dunaliella* grows faster at lower salinity (less than roughly 22 percent (w/v). NaCl) saturation) but has the maximum beta-carotene concentration at greater salinity. (about 33 % of NaCl).



Figure 2.5: *Dunaliella* under microscope (Rappe, 2019).

Although *Dunaliella* grows more rapidly at low salinity, there is the potential for contamination by protozoa. Ben-Amotz studied numerous ways of producing *Dunaliella* (Borowitzka & Vonshak, 2017). A nitrogen deficiency, an increase in salt content, and high sun radiation are characteristics of intensive culture environments that encourage the growth of *Dunaliella* biomass and the production of carotene. (Borowitzka & Vonshak, 2017). Typically, *Dunaliella* is harvested using centrifugation, flocculation, or a process that utilizes the cell membrane's hydrophobic characteristics (Borowitzka & Larkum, 1987). Once harvested, biomass can be dried using a spray or tumble dryer, or carotene can be directly extracted using heated oil or other solvents. The biological composition of *Dunaliella* consists of 49–57% protein, 4–32% carbs, and 6–8% lipids. (Zhu et al., 2014).

2.2.4 Haematococcus

The unicellular Chlorophyte microalga *Haematococcus pluvialis* (*H. pluvialis*) is found in freshwater across the globe (Figure 2.6). It is well known that this species accumulates astaxanthin of up to 2–3 percent by dry weight under stressful conditions. the most significant commercial producer of astaxanthin is *H. pluvialis* (Ambati et al., 2014). *H. pluvialis* is grown in various

photobioreactors, including closed photobioreactors, open raceway ponds, and photobioreactors of various types (based on growth conditions) (Kang et al., 2010; Wang et al., 2013).



Figure 2.6: *Haematococcus* sp. under microscope (Fox, 2022).

The majority of *H. Pluvialis* photoautotrophic culture occurs in open raceway ponds or enclosed photobioreactors, where algae thrive. For commercial production, a two-stage cultivation approach is generally used (Shah et al., 2016). Environmental variables such as light, pH, temperature, salt content, and nutritional stress affect astaxanthin accumulation (Shah et al., 2016).

2.2.5 *Schizochytrium* sp.

Schizochytrium sp. species were formerly classified as fungi due to their heterotrophic nature and similarity to Chytrids. However, this organism was categorized as heterokont algae in the Thraustochytrids by using contemporary molecular biology methods (Cavalier-Smith, 1999). *Schizochytrium* sp. microalgae may be regarded as a component of the human food chain because it is consumed by fish and other marine organisms. The search for alternatives to fish oil has risen due to the increased focus on the benefits of including omega -3 fatty acids in both human and animal diets. As a result of the growing need for omega 3 fatty acids in human diets, the hunt for alternatives to fish oil has grown significantly. *Schizochytrium* sp. is one of a number of single-celled organisms that generate polyunsaturated fatty acid (PUFA)-rich oils, including EPA and DHA (Figure 2.7) (Vardar, 2016).




Figure 2.7: *Schizochytrium* sp. cells under microscope (Vardar, 2016).

Manufacturing *Schizochytrium* sp. has several advantages, such as improved oxidative stability, reduced purification costs, production from natural and sustainable ingredients, and consistent product quality. (Ratlidge, 2004). *Schizochytrium* sp. is one of the most widely used commercial strains owing to its PUFA synthase complex's O₂-independent activity (Jakobsen et al., 2008). Under oxygen-limiting circumstances, the lipid content of *Schizochytrium* sp. may rise, which may impact the activity of the PUFA synthase enzyme complex. The changing dissolved oxygen level in a shaking flask scale was also examined, and DHA content was found to be elevated, according to a two-stage growth experiment with *Schizochytrium* sp. (Chi et al., 2009).

2.3 COMPOSITION OF MICROALGAL BIOMASS

Microalgae are very adaptable to changing environmental circumstances, displaying exceptional flexibility and adaptability. As a consequence of their defensive and adaptation systems, they create a wide range of biosynthetic chemicals or molecules with distinct features (Guschina & Harwood, 2006). There is significant potential for some of these chemicals to be used as precursors or raw materials for value-added bioproducts in various industries. Components of microalgae (such as proteins, carbohydrates, and lipids) enable them to adapt to fluctuating environmental circumstances for growth (Mooij et al., 2016; Markou et al., 2015). Typical microalgae nutritional products include powders, pills, and capsules that provide between 12 to 35 % protein, 7.2 to 23 % fat, and 4.6 to 23 % carbohydrates (Borowitzka, 2013).

Microalgal biomass content, especially lipid production, is significantly affected by key abiotic variables such as light intensity, medium composition, temperature, pH, salt content, and nutrient restriction (especially nitrogen restriction) (Pal et al., 2011). The effects of these traits are species-specific; therefore, ideal conditions for one microalgae species may not necessarily be advantageous for another microalgae species (Robertson et al., 2013). Understanding how these parameters interact with metabolic pathways in different microalgae species is therefore crucial for making well-informed decisions.

2.3.1 Carbohydrates

Microalgae are the principal producers of biomass, such as carbohydrates, proteins, and lipids throughout the planet by photosynthesizing light energy and carbon dioxide (CO₂) (Becker, 2007). Microalgae synthesize carbohydrates for two primary functions: as a structural component of the cell wall and as a storage component inside the cell (Cheng et al., 2017; Schulze et al., 2016). As a storage molecule, carbohydrates supply the microalgae with the energy required for their metabolism and enable them to temporarily live in the dark when necessary (Fimbres-Olivarria et al., 2016). Carbohydrate concentration in biomass is dependent on microalgal species and growing circumstances (Patel et al., 2014). Certain microalgal species, such as *Porphyridium cruentum* (40–57%) and *Spirogyra* sp. (33–64%), contain a great deal of carbohydrates (Mühlroth et al., 2013; Dai et al., 2020). Microalgal polysaccharides are used in the generation of biofuels, particularly biohydrogen. To improve biofuel production, it is required to combine circumstances with a high carbohydrate content with microalgae's ability to generate substantial volumes of biomass. Therefore, culture conditions can be changed to increase the growth rate and carbohydrate content of microalgae (Blockx et al., 2018; Chng et al., 2017).

Microalgal carbohydrate concentration is primarily influenced by nutritional content, stress conditions, light intensity, and temperature. The technique of lowering or raising the quantity of nutrients is regarded as a cost-effective method for producing microalgae high in carbohydrates (Corrêa et al., 2019). This is feasible because it's easy to regulate the medium's nutritional content. (Corrêa et al., 2019; Gerchman et al., 2017).

2.3.2 Protein

Many species of microalgae have significantly variable protein compositions. For example, *Chaetoceros gracilis* has 12% protein, while *Arthrospira platensis* has between 53-70% protein (Wang et al., 2021). The protein content in the microalgal biomass is affected by many factors such as genetic characteristics, light intensity, temperature, pH, CO₂ content, medium content, and the culture system. These factors, affect the protein content of different microalgae species differently. The amount of protein obtained from microalgae can also vary according to the extraction method. Under the ideal circumstances of biomass concentration, ionic strength, temperature, pH, and extraction time, microalgae proteins can typically be extracted from acidic, aqueous, and/or alkaline media. After centrifuging the medium, proteins may be recovered using chromatography, ultrafiltration, and separation methods like precipitation. The isolation of crude protein concentrates using this method is straightforward, although there is no universally applicable pH value for the extraction of microalgae protein. For instance, *Chlorella vulgaris* and *Spirulina platensis* reported the maximum protein solubility achieved at pH 11 and pH 12, respectively (Ursu et al., 2014), but a higher value of protein was isolated from *Nannochloropsis* sp. up to pH 13.(Gerde et al., 2013).

2.3.3 Lipids

Microalgae cells use both organic and inorganic carbon sources for lipid production. The lipid content of microalgae generally ranges from 10-60% of dry biomass. However, under certain conditions, the microalgal lipid content can even reach 80%. Microalgal lipids can be divided into two different classes based on their carbon number. Polyunsaturated fatty acids (PUFA) with more than 20 carbons especially docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) are frequently utilized as health food supplements, while fatty acids with 14–20 carbons are mostly employed for the manufacturing of biodiesel. It is known that the DHA range of *Schizochytrium* sp. ranges from 20% to 40% of the total amount of fatty acids.

Under stressful circumstances like nutrition restriction and low temperature, microalgae can synthesize significant levels of lipids. For example, lipid accumulation in *Dunaliella* sp. and *Chlorella vulgaris* increased significantly in high-salt media, reaching 70% and 21%, respectively.

In another study, observations revealed that the lipid ratio of *Schizochytrium* sp. microalgae increased 4-fold in a nitrogen-limited environment.

Industrial output is supported by the capacity of green algae to accumulate carotenoids, cell proliferation, and lipid content (Aguilar-Rosas et al., 2013). Microalgal lipid amounts can be increased by overexpression of key enzymes involved in lipid biosynthesis. Lipid synthesis may be improved by manipulating DGAT levels. For example, in *Arabidopsis thaliana*, DGAT overexpression resulted in a 10-70 % increase in lipid content (Jako et al., 2001). Another study found that the DGTT2 gene could be successfully transplanted into *Arabidopsis* after being extracted from *Chlamydomonas reinhardtii*. The quantity of lipids in transgenic plants increased by about ten times, and the newly generated lipids included exceptionally long-chain fatty acids. Sometimes, the overexpression of enzymes in fatty acid metabolism is not effective at all in increasing the total fat ratio. ACCase is an enzyme widely acknowledged as the building block of fatty acids, and hence it is critical to the metabolism of lipids (Dunahay et al., 1996). Excessive overexpression of ACCase in diatoms and other plants was ineffective in increasing lipid synthesis. Even in the case of the diatom *Cyclotella cryptica*, a study in 1996 found that overexpressing ACCase did not affect the diatom's ability to produce fat.

2.4 LONG CHAIN POLYUNSATURATED FATTY ACIDS (PUFA)

There are several metabolic processes in which fatty acids play an essential role (Robertson et al., 2013). Polyunsaturated fatty acids (PUFAs) are important for various functions such as enzyme interactions, membrane fluidity, and lipid peroxidation (Borowitzka & Vonshak, 2017). An important role of polyunsaturated fatty acids in cell signaling and inflammatory processes has been demonstrated in many studies (Mozaffarian and Wu, 2011). Some disorders, such as coronary artery disease, cardiac arrhythmias, diabetes, hyperlipidaemia, and atherogenesis, can be exacerbated by poor dietary habits. (Surette, 2008). Supplementing LC-PUFAs to diet has been demonstrated to lower the chance of getting Alzheimer's and serious depression. Lipids, namely triacylglycerol (TAGs), are regarded as universal energy storage molecules and play an essential role in cellular metabolism (Xin et al., 2019). TAGs may be exploited as nutritional supplements or biofuels, depending on their fatty acid makeup (number of carbons, chain length, and degree of

unsaturation). Polyunsaturated fatty acids (PUFAs) are necessary components of the human diet and are crucial for the maintenance of proper growth and health (Petrie et al., 2010).

Triacylglycerol (TAG) plays an important role in cellular metabolism and membrane structure. TAGs are also known universally as energy storage molecules (Xin et al., 2019). Depending on the fatty acid in the structure of TAGs, it can also be used as a nutritional supplement, mostly as a biofuel. PUFAs (polyunsaturated fatty acids), on the other hand, contain important components of the human diet with their long chain and unsaturated carbon structure and are very important for healthy growth and protection of health (Petrie et al., 2010). (LC-PUFAs) are classified according to the carbon where the first double bond is closest to the methyl group. Omega-3 fatty acids are called " ω -3 fatty acids" or "n-3 fatty acids" because they contain a double bond three atoms away from the terminal methyl group in their chemical structure. Alpha-linolenic acid (ALA), which is found in vegetable oils, EPA, and DHA, which are both often found in marine fish oils, are three different forms of omega-3 fatty acids that are important in human physiology.

Omega-6 fatty acids are called " ω -6 fatty acids" or "n-6 fatty acids" because they contain a double bond six atoms away from the terminal methyl group in their chemical structure. Fatty acids with omega-6 bonds cannot be produced in the human body, so they are called essential (essential) fatty acids. Examples of omega-6 fatty acids are linoleic acid (18:2, n-6) and arachidonic acid (20:4, n-6). Deficiency in essential fatty acids can cause diseases, especially since they are found in the brain.

Omega-9 fatty acids are called " ω -9 fatty acids" or "n-9 fatty acids" because they contain a double bond nine atoms away from the terminal methyl group in their chemical structure. The two most well-known types of omega 9, which are mostly found in products such as olive oil, are oleic acid and meadow acid. Unlike omega-3 and omega-6, it does not show essential fatty acid properties.

2.4.1 Omega-3

Alpha-linolenic acid (ALA), Eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are omega-3 polyunsaturated fatty acids known to have beneficial effects on human health and nutrition. The omega-3 fatty acids important in the human diet are alpha-linolenic acid (18:3, ALA), (20:5, EPA eicosapentaenoic acid), and (22:6, DHA docosahexaenoic acid). Polyunsaturated fatty acids, which have beneficial effects on human health and nutrition, have

been the subject of study for many years. It guards against high blood pressure, lowers triglyceride levels, raises HDL, or "good cholesterol," which helps lower the risk of heart attack and stroke, and preserves cardiovascular health. It has an anti-inflammatory effect and has potential effects in the control of cell proliferation and the fight against tumors. It is known that it contributes to the reduction of symptoms of some neurological and psychiatric diseases and to the prevention of age-related vision loss. Omega-3 is also very important for the brain development, the development of the central nervous system, and the bodily development of babies in the womb. (Green et al., 2008; Chin et al., 1992; Mattson & Grundy, 1985).

Some studies have shown that DHA and EPA, two important omega-3 PUFAs, have antimicrobial activity (Sun et al., 2016). The antibacterial effects of EPA and DHA that appear to target the membrane of the bacterial cell include disruption of the electron transport chain and decoupling of oxygen-dependent phosphorylation, enzyme inhibition, cell necrosis, depletion of nutritional intake, and autooxidation (Sun et al., 2016; Desbois & Smith, 2010). Due to the amphipathic nature of EPA and DHA, they follow an action path that can make contact with the cell membrane and cause holes in the membrane. Both gram-negative and gram-positive types of bacteria are inhibited by EPA and DHA, while gram-positive bacteria are more successfully neutralized by EPA and DHA. (Desbois & Smith, 2010; J. Sun & Song, 2011).

2.4.1.1 Eicosapentaenoic acid (EPA)

Eicosapentaenoic acid is an omega-3 fatty acid that shown as 20:5 because it has twenty carbon chains and five cis double bonds in its chemical structure (Fig 2.8). EPA, an essential omega-3 fatty acid, is critical for the circulatory system in animals because it inhibits platelet aggregation (Parka & Harris, 2002). Additionally, the availability of the essential EPA dosage throughout the human body aids in inflammatory management and improves the immune system (Calder, 2013). Additionally, EPA is critical in preventing heart disease and reducing chronic and inflammatory processes (Swanson et al., 2012). DHA and EPA also play a significant role in the body's defensive system by lowering inflammation and eliminating reactive oxygen species at the cellular level. Since our body cannot produce enough EPA and DHA according to its own needs, it should be supplemented with nutrients in appropriate proportions. (Nichols et al., 2010).

2.4.1.2 Docosahexaenoic acid (DHA)

An omega-3 fatty acid called docosahexaenoic acid serves as the main structural element of the human brain, skin, cerebral cortex, and retina. Due to its twenty-two carbon chains and six cis double bonds in its chemical structure, it is represented as 22:6 (Figure 2.8). Human diseases like cancer, Alzheimer's disease, cardiovascular disease, and schizophrenia can all be prevented with DHA (Lee & Lip, 2003).

Animals, like humans, receive omega-3 fatty acids from dietary sources (Certik & Shimizu, 1999). Coldwater marine fish are the best-known commercial source of these fatty acids worldwide. Due to its high DHA content, its commercial value has increased during the last several decades. However, it is known that animals (fish) lack the enzymes required to produce DHA, which should be supplemented externally. In contrast, microalgae are the primary natural producers of omega-3 and omega-6 fatty acids (Nichols et al., 2014). Some microalgae species living in the aquatic environment are rich in DHA. *Schizochytrium* sp. microalgae have become more attractive as a dietary supplement due to their high DHA content. (Apt & Behrens, 1999). By directly ingesting microalgae and/or indirectly via other tiny fish and zooplanktons that consume microalgae, most fish may retain significant levels of DHA and EPA (Peltomaa et al., 2018; Nichols et al., 2014).




Figure 2.8: Molecular structures of DHA and EPA fatty acids

2.5 ANTIMICROBIAL ACTIVITY OF MICROALGAE

Although microalgae have been used for medicinal purposes for a long time, systematic screening for biologically active components began in the mid-1900s. However, microalgae have been the subject of intensive study aimed at discovering new chemicals that may lead to therapeutically effective medicines in the last decade (Mendes et al., 2003; Cardozo et al., 2007). Microalgae contain many extracts and/or extracellular products with proven antibacterial, antiprotozoal,

antifungal, and antiplasmodial properties (Kellam & Walker, 1989; Ghasemi et al., 2004). The antibacterial effect of microalgae has been associated with many chemical classes such as terpenes, phenolics, and amines. An organic extract of *Anabaena virabilis*, a blue-green algae species, obtained with hexane and methanol has been shown to be effective on vancomycin-resistant *S. aureus* strains (Mayer & Hamann, 2005; Cardozo et al., 2007). The bioactive components of marine algae species related to different divisions, such as Chlorophyta and Rhodophyta, were extracted using solvents such as acetone, ethanol, and methanol-toluene. The antibacterial and antifungal activities of the related extracts were evaluated. Studies in Rhodophyta spp. species such as *Laurencia okamurai*, *Grateloupia*, and *Plocamium telfairiae* have shown that these species have a broad spectrum of antibacterial activity. Some strains were found to show very strong antimicrobial activity against test organisms. Among these test organisms, it was found to be particularly sensitive to extracts of *Pseudomonas solancearum*, and *Penicilium citrinum* marine algae, and Rhodophyta spp. In these results, it was defined that marine algae and Rhodophyta spp. extracts have considerable potential for application in the creation of novel medications (Yi et al., 2001).

Another study examined the antibacterial activity of six different microalgae extracts of Chlorophyta and Rhodophyta using several solvents, such as ethanol, hexane, and chloroform. According to the findings, *Amansia multifida* had significant antibacterial activity against gram-negative pathogens including *Salmonella typhi*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*., *Salmonella choleraesuis*, and *E. aerogenes* and gram-positive bacteria such as *Bacillus subtilis* and *S. aureus* (Lima-Filho et al., 2002). Similar findings were discovered by Bhagavathy et al. (2011) who assessed the presence of the phytochemicals and antibacterial properties of *Chlorococcum humicola*. The screening of *C. humicola* extracts revealed antibacterial efficacy against pathogenic strains including *K. pneumoniae*, *B. subtilis*, *P. aeruginosa*, *V. cholerae*, *Salmonella typhi*, *S. aureus*, *Aspergillus flavus*, *Candida albicans*, *Aspergillus niger*, and *E. coli*.

Interestingly, several studies indicate that algae antimicrobial compounds undergo seasonal and growth-related changes. For example, Hornsey and Hide (Bourgougnon, 2011) showed that seasonal fluctuations in antibacterial content in examined algae occurred in four major patterns:

- a. The *Polysiphonia* kind, which produces antibiotics continuously throughout the year.
- b. *Laurencia Pinnata*, *Chondrus crispus*, and *U. lactuca* also include the *Laminaria* type, which has the highest antibiotic production, especially in winter.
- c. The *Dictyota* type has an activity peak mostly in the summer and is found in *Ascophyllum nodosum* and *Dilsea carnososa*.
- d. The *Codium* type, characterized by a springtime activity peak, as shown by *Halidrys siliquosa*.

Today, new generation techniques such as environmentally friendly new extraction methods and modern analytical devices are used to screen new products and determine the biological activity of microorganism extracts. By using new generation devices such as GC-MS and HPLC-DAD, antioxidant and antimicrobial compounds that may show antimicrobial activity against pathogenic species such as *E. coli*, *S. aureus*, and *C. albicans* are being screened (Plaza et al., 2010).

2.5.1 Antibacterial Activity

The resistance of harmful bacteria to antibacterial drugs has increased significantly over the past decade. Therefore, new antimicrobial treatment options for diseases caused by resistant microorganisms have gained considerable interest in addition to our understanding of the process of antimicrobial resistance. Selected compounds from various microalgae and their antibacterial properties are shown in Table 2.1.

Table 2.1: Antibacterial properties of compounds from microalgae (Amaro et al., 2011).

Microalga	Active compound	Target microorganism
<i>Phaeodactylum tricornutum</i>	Eicosapentaenoic acid	MRSA, <i>Listonella anguillarum</i> , <i>Lactococcus garvieae</i> , <i>Vibrio</i> spp.
<i>Haematococcus pluvialis</i>	Short chain fatty acids	-
	Short-chain fatty acids (butanoic acid and methyl lactate)	<i>Escherichia coli</i> , <i>Staphylococcus aureus</i>
<i>Skeletonema costatum</i>	Unsaturated, saturated long chain fatty acids	<i>Vibrio</i> spp.
<i>Euglena viridis</i>	Organic extracts	<i>Pseudomonas</i> , <i>Aeromonas</i> , <i>Edwardsiella</i> , <i>Vibrio</i> , <i>E. coli</i>
<i>S. costatum</i>	Extra-metabolites	<i>Listeria monocytogenes</i>
<i>Staurostrum gracile</i>	Methanolic extracts	-
<i>Pleurastrum terrestre</i>		
<i>Dictyosphaerium pulchellum</i>		
<i>Klebsormidium crenulatum</i>		
<i>Chlorococcum</i> sp.	Aqueous extract	-
<i>Chlorococcum</i> HS-101	α -Linolenic acid	-
<i>Chlorokybus atmophyticus</i>	Acetone extract	-
<i>Chlamydomonas reinhardtii</i>	Methanolic and hexanolic extracts	<i>S. aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Bacillus subtilis</i> , <i>E. coli</i> , <i>Salmonella typhi</i>
<i>Chlorella vulgaris</i>		

Antibiotic resistance in bacteria has been the focus of research for the following reasons: (i) The majority of community-acquired and hospital-acquired illnesses are caused by bacteria. (ii) An increase in the number and dose of antibacterial agents provides a wider variety of resistance. (iii)

The capacity to transfer bacterial resistance makes it possible to delve deeper into the processes underlying resistance formation (Amaro et al., 2011).

Meanwhile, research aimed at discovering the active principles of microalgal antibacterials has gained popularity. The first antibacterial chemical, chlorellin, was discovered in *Chlorella* microalgae and was found to be responsible for its inhibitory effect against Gr⁺ and Gr⁻ bacteria (Pratt et al., 2022). Eicosapentaenoic acid, a compound produced by *P. tricornutum* diatoms, has been shown to be responsible for its antimicrobial activity against Gr⁻ and Gr⁺ bacteria (including MRSA). However, two different antibacterial properties of free fatty acids (C16:1 n-7; palmitoleic acid) and hexadecatrienoic acid (HTA; C16:3 n-4) produced by *P. tricornutum* were also discovered. It has been shown to inhibit Gram-positive bacteria with HTA as well as the Gram-negative marine pathogen *Listonella anguillarum*. (Desbois et al., 2008). While free fatty acids can act on different cellular targets, cell membranes are the most plausible in terms of antibacterial activity. Because damage to the cell membrane will likely lead to disruption of cell integrity and reduced nutrient absorption in addition to limiting cellular respiration. For example, chemicals produced by *Scenedesmus costatum* and partly isolated from its organic extract displayed antibacterial action against aquaculture bacteria due to the presence of fatty acids with chain lengths greater than 10 carbon atoms, which trigger the lysis of bacterial protoplasts (Desbois et al., 2009). It has been known for some time that fatty acids have the potential to inhibit bacterial growth and survival, but new structure-function connection studies reveal that this ability depends on both the degree of unsaturation and the length of the chain (Benkendorff et al., 2005).

Among microalgal-derived oxylipins, polyunsaturated aldehydes stand out for their antibacterial properties. These chemicals are produced by some diatoms, including *Thalassiosira rotula* and *S. costatum*. There are studies in which (C20:4 n-3) arachidonic acid shows effective activity against significant human pathogens, such as *Haemophilus influenza*. In this study, the MIC values for MRSA and *H. influenza* were found to be 1.9 µg/mL and 7.8, respectively. It also showed that it impaired the growth of gram-negative bacteria such as *L. anguillarum*, *Photobacterium phosphoreum*, *Aeromonas hydrophila*, and *Alteromonas haloplankti*, and gram-positive marine bacteria such as *Micrococcus luteus* and *Planococcus citreus* (Smith et al., 2010).

2.5.2 Antiviral Activity

In recent years, various infectious disorders caused by several viruses have developed (and re-emerged). Despite the fact that a number of antiviral medications have been created, drug-resistant mutations are continually occurring, necessitating the development of novel antiviral active principles. As a result, microalgae have attracted considerable interest as possible sources of antiviral medicines. Selected compounds from various microalgae and their antiviral properties are shown in Table 2.2.

Table 2.2: Antiviral properties of compounds from microalgae (Amaro et al.,2011).

Microalga	Active compound	Mechanism of action	Target virus
<i>Navicula directa</i>	Polysaccharide	Inhibition of hyaluronidase	HSV1 & 2, Influenza A virus
<i>Gyrodinium impudicum</i>	p-KG03 exopolysaccharide	Inhibition (or slowing down) of cytopathic effect	Encephalomyocarditis virus
<i>Dunaliella primolecta</i>	Pheophorbide α -, β -like compounds	Inhibition of cytopathic effect	HSV1
<i>Chlorella autotrophica</i>	Sulfated polysaccharides	Replication inhibition C. <i>autotrophica</i> : 47.4-67.4 % <i>Ellipsoidon</i> sp.: up to 44 %	VHSV, ASFV
<i>Ellipsoidon</i> sp.			
Cryptomonads	Allophycocyanin	Inhibition of cytopathic effect, delay in synthesis of viral RNA	Enterovirus 71
<i>Cochlodinium polykrikoide</i>	Extracellular sulfated polysaccharides	Inhibition of cytopathic effect 1	Influenza virus A & B, RSV A & B, HSV-1

Antiviral activity may occur during a single or many phases of viral growth. Phase I is concerned with cell adsorption and invasion; Phase II, or the eclipse phase, is distinguished by the cell's compulsion to produce numerous copies of the virus; and Phase III, which is concerned with viral

particle maturation and release. Each antiviral compound can act at different stages. For example, the anti-HSV effect of an antiviral agent, acyclovir, occurs in phase II, while the anti-HSV component produced from *Dunaliella sp* inhibits viral function in phase I (Elion et al., 1977). It has been suggested that sulfated exopolysaccharides from marine microalgae interact with phase I of some enveloped viruses (Batinić & Robey, 1992). They provide a competitive advantage against viruses such as HSV and HIV due to their wide antiviral range (Veettil et al., 2022). Apparently, their inhibitory effect is due to their interaction with positive charges on the virus surface, preventing the virus from invading host cells (Ehresmann et al., 1979). The administration of such compounds appears to be an especially promising antiviral therapeutic option because their pleiotropic mode of action makes the development of resistance mutants less likely than with other antiviral drugs that demonstrate just a single target throughout the viral life cycle. Analyses of antiviral substances produced by *D. primolecta* were performed by NMR and MS. As a result of the study, the pheophorbide-like compounds that had the proton at position 21 substituted by a hydroxyl group that may account for their exceptional antiviral activity were detected (Ohta et al., 1998).

2.5.3 Antifungal Activity

Until the last few years, studies on the development of resistance to antifungal agents had not received as much attention as the study of antibacterial resistance. (Anaissie et al., 1989; Wey et al., 1988). This is because fungi are not seen as being just as pathogenic as bacteria. While the death rate from fungal infections was stable in the mid-1900s, it has increased in recent years. During this rise in fungal infections, researchers have begun to look for new and effective antifungal drugs to fight the growing number of fungal infections. Selected compounds from various microalgae and their antifungal properties are shown in Table 2.3.

Table 2.3: Antifungal properties of compounds from microalgae (Amaro et al.,2011)

Microalga	Active compound	Target microorganism
<i>Chlamydomonas reinhardtii</i>	Methanolic extracts	<i>Candida kefir</i> , <i>Aspergillus niger</i> , <i>Aspergillus fumigatus</i>
<i>Chlorella vulgaris</i>		
<i>Oocystis</i> sp.		
<i>Scenedesmus obliquus</i>		
<i>Amphidinium</i> sp.	Karatungiols	<i>A. niger</i> , <i>Trichomonas foetus</i>
<i>Goniodoma pseudogoniaulax</i>	Goniodomin A	-
<i>Gambierdiscus toxicus</i>	Polyether compounds (gambieric acids A and B)	-
<i>Prorocentrum lima</i>	Polyether compounds	-
<i>Dinophysis fortii</i>		-
<i>Haematococcus pluvialis</i>	Butanoic acid and methyl lactate	<i>Candida albicans</i>
<i>Chlorella pyrenoidosa</i> , <i>Scenedesmus quadricauda</i>	-	<i>A. niger</i> , <i>Aspergillus flavus</i> , <i>Penicillium herquei</i> , <i>Fusarium moniliforme</i> , <i>Helminthosporium</i> sp., <i>Alternaria brassicae</i> , <i>Saccharomyces cerevisiae</i> , <i>C. albicans</i>

Katircioğlu et al. (Katircioğlu et al., 2006) studied the antifungal properties of ten microalgae strains isolated from different fresh water sources in Turkey against *S. cerevisiae*, *Candida tropicalis*, and *C. albicans*. According to the results they obtained, *Oscillatoria* sp. and *Chlorococcus* sp. strains showed the best antifungal activity.

In another study by Santoyo et al. (Santoyo et al., 2009), extracts of *H. pluvialis* obtained with pressurized liquid and ethanol were tested on *C. albicans*. It has been suggested that two compounds with antifungal effects are butanoic acid and methyl lactate.

Abedin and Taha (Abedin & Taha, 2008) investigated the antifungal activities of *S. quadricauda* and *C. pyrenoidosa* microalgae extracts on eight different fungi. According to their results, all the extracts showed positive effects on almost all fungi. The results obtained from this study are in line with similar studies done before.

2.6 ALGAL FATTY ACIDS AS ANTIMICROBIAL COMPOUND

Free fatty acids are the lipids in human skin that are thought to be the most active. The most important of these fatty acids are lauric acid (C12:0), palmitic acid (C16:0), myristic acid (C14:0), and C18:1n-7 (cis-8-24 octadecenoic acid) (Stabili et al., 2012). Algae species such as *Haptophyta* and *Cryptomonas* are known to contain significant amounts of polyunsaturated fatty acids (PUFAs) and saturated fatty acids (SFAs). These fatty acids are important for human health and also because they cannot be produced by humans and must be taken from outside (Ryckebosch et al., 2014). A list of bioactive molecules isolated from different algal species is summarized in Table 2.4.

Table 2.4: A list of bioactive molecules isolated from different algal species (Parsaeimehr et al., 2016).

Bioactive Molecule	Algal Source	Mode of Action	Target
3-Formylindole	<i>Botryocladia leptopoda</i>	Antiviral and anticancer activity	<i>Vaccinia</i> virus
Polyhydroxylated	<i>Fucus vesiculosus</i>	Bactericidal	Gram-positive and Gram-negative bacteria
Fucophlorethol	<i>Ascophyllum nodosum</i>	Antitumor activity	HT-29 and HCT-116 human colon cancer
Fucoidan	<i>Symphyocladia latiuscula</i>	Bactericidal	<i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , and <i>Pseudomonas aeruginosa</i>

Table 2.4 continued: A list of bioactive molecules isolated from different algal species
(Parsaeimehr et al.,2016).

Bioactive Molecule	Algal Source	Mode of Action	Target
Sulfated galactans	<i>Botryocladia occidentalis</i>	Immunostimulatory activity, antiviral and anticoagulant activity	Herpes simplex virus, human metapneumovirus
Phycarine	<i>Laminaria digitata</i>	Antitumor activity	Stimulator of both hormonal and cellular immunity, and stimulator of phagocytosis by peripheral blood cells
Loliolide	<i>Undaria pinnatifida</i>	Antioxidant, bactericidal, and antitumor activity	<i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>S. aureus</i> , and <i>Neisseria gonorrhoeae</i>
Bisebromoamide	<i>Lyngbya sp.</i>	Anticancer activity	Renal cell carcinoma
p-KG03 exopolysaccharide	<i>Gyrodinium impudicum</i>	Antifungal HSV-1 Antiprotozoan Antiviral	HSV-1, <i>Aspergillus niger</i> , <i>Trichomonas foetus</i> <i>Encephalomyocarditis virus</i>
Allophycocyanin	<i>Spirulina platensis</i>	Antiviral	<i>Enterovirus 71</i>
Calothrixin-A	<i>Calothrix sp.</i>	Antimalarial and anticancerous	Human myeloid leukemia K562 cells
Sulfoquinovosyldiacylglycerol, KM043	<i>Gigartina tenella</i>	Antibacterial	Inhibitor of eukaryotic DNA polymerases and HIV reverse transcriptase type 1
Deschloroelatol	<i>Laurencia rigida</i>	Antifungal	<i>Mycotypha microspora</i>

Many microalgae species have palmitic acid in their fatty acid composition, which is saturated fatty acids and has demonstrated antiviral effects. Santoyo et al. (Santoyo et al., 2012) reported that palmitic acid isolated from *Haematococcus pluvialis* and *Dunaliella salina* showed anti-HSV-1 activity.

The *Polysiphonia virgata*'s extract which was obtained by dichloromethane, showed antimicrobial activity against *Mycobacterium tuberculosis* and *M. smegmatis*. The results obtained show that the fatty acids isolated from *Polysiphonia virgata*, especially oleic acid, lauric acid, linoleic acid, and myristic acid, have antimicrobial properties (Saravanakumar et al., 2008).

The antimicrobial activity of *Dunaliella salina*, which is a halophilic green microalga and widely grown commercially for the production of β -carotene, on microorganisms such as *E. coli*, *C. albicans*, *S. aureus* was investigated. When *Dunaliella salina* extracts were tested with the GC-MS device, 15 different antimicrobial compounds were reported, including β -cyclocitral, phytol, fatty acid, and β -ionone (Herrero et al., 2006).

In another study, the antimicrobial activity of extracts of *Dunaliella primolekta* and *Chlorococcum* microalgae using methanol was tested on methicillin-resistant *S. aureus*. As a result, it was determined that the extracts showed antimicrobial activity against the resistant organism. In addition, it was observed that the antibiotic production of *Dunaliella primolekta* and *Chlorococcum* increased 1.8-2.3 times in the developed BG-11 medium (Ohta et al., 1995).

Rodríguez-Meizoso et al. (2010) studied the antimicrobial activity of *Haematococcus pluvialis*, a freshwater microalga that produces high levels of astaxanthin. They analyzed the extracts of *Haematococcus pluvialis* using analytical techniques such as HPLC-DAD, HPLC-MS, and GC. According to the results they obtained, it was determined that short-chain fatty acids were responsible for the antimicrobial activity.

When the studies are examined, it is possible to say that the hydroxyl groups of fatty acids have antimicrobial properties. Because it has been determined that antimicrobial properties of methylated fatty acids that do not contain hydroxyl groups are very low or none at all. Studies have shown that the antimicrobial activities of medium and long chain unsaturated fatty acids are higher than those of saturated fatty acids of the same length (Desbois et al., 2008). In general, there is a very important correlation between the number of the double bonds in the carbon chains of

unsaturated fatty acids and the antibacterial activity of fatty acids (Feldlaufe et al., 1993). On the other hand, it is known that saturated fatty acids with a length of 10-12 carbons also show very strong antimicrobial activity (Wille et al., 2003).

When the data obtained from the studies are examined, it has been revealed that fatty acids show antimicrobial activity behaviour on microorganisms in many different ways, such as disruption of the electron transport chain, cell lysis, peroxidation and autoxidation, and deterioration of food intake (Desbois et al., 2010).

2.7 ANTIBIOFILM ACTIVITY

The term "biofilm" refers to the microbial population that is adherent to abiotic or biotic surfaces. The process of forming a biofilm involves several steps, including the "first adhesion" stage, in which bacteria adhere to a surface, the "micro-colony formation" stage, in which bacteria divide to produce extracellular polymeric molecules (EPS) that position the cells and increase adhesion, the "biofilm maturation" stage, in which the EPS matrix forms a three-dimensional structure in the form of a multifunctional and protective skeleton and the "splitting" (also called dispersal) stages, in which bacteria can break away from the biofilm and recolonize new areas. There are two strategies for dealing with biofilms: preventing biofilm formation or eliminating the growing biofilm. Biofilm-forming bacteria differ from planktonic bacteria in that they have immune responses and extreme tolerance to antimicrobial therapy. Therefore, biofilm formation requires higher doses of antibiotics and longer duration of therapy in the treatment of chronic infections (Wannigama et al., 2020). The genetic variety of organisms that create biofilms demonstrates that biofilms are a primitive form of microbial human life (Uzundağ, 2020). Van Leeuwenhoek made the first discovery of biofilm on the tooth surface (Between & Concentration, 1940). Biofilms are believed to be a collection of adherent microorganisms (Hall-Stoodley & Stoodley, 2009). Bacterial cells are protected by biofilms from severe environmental conditions, and as a result, they are particularly resistant to antibiotics.

In industrial, ecological, and medicinal environments, biofilms are exceedingly dangerous. For instance, biofilms on the surface of medical equipment such as implants or catheters commonly result in infections that are difficult to cure. On the other hand, infections have been linked to the production of biofilms on the surfaces of the human urinary system, skin, and teeth (Romeo, 2008).

Dental plaque biofilms contain more than one species, and the disease-causing capacity of the community is often affected by the structure of the biofilm. (Kreth et al., 2008). *E. coli* has been warned that it is capable of forming biofilms in the human gut, posing a substantial danger to human health (Van Houdt & Michiels, 2005).

While the antimicrobial properties of many compounds produced by microalgae and blue-green algae have been examined, very few of them have been examined in terms of their antibiofilm properties. Selected compounds from various microalgae and their antibiofilm properties are shown in Table 2.5.

Table 2.5: Selected compounds from various microalgae and their antibiofilm properties (Nag et al.,2022)

Name of the active compound/extract	Algae	Activity
Galactoglycerolipids	<i>Sargassum muticum</i>	Exhibits antibiofilm activity
Mahorone, 5-bromomahorone	<i>Asparagopsis taxiformis</i>	Helps in the eradication of biofilm
1,1,3,3-Tetrabromo-2-heptanone	<i>Bonnemaisonia hamifera</i>	Helps in the eradication of biofilm
Linear alkadienes	<i>Botryococcus braunii</i>	Helps in the eradication of biofilm and are also responsible for the process of phycoremediation
Neophytadiene, tocopherol, palmitoleic acid, phycocyanin	<i>Spirulina platensis</i>	Acts as potent antimicrobial and antibiofilm agent
Astaxanthin	<i>Chlorella zofingiensis</i>	Has effective antibiofilm properties
Palmitoleic acid C-phycocyanin, oleic acid, PUFAs (n-3) fatty acids, phytol, phenolic acids, neophytadiene linolenic acid	<i>Spirulina platensis</i>	Helps in the eradication of biofilm

Methanol and ethanol extracts	<i>Ulva fasciata</i>	Helps in the eradication of biofilm
Chromanols	<i>Sargassum horneri</i>	Has effective antibiofilm properties
15,16-epoxythysiferol B; 15,16-epoxythysiferol A; 28-hydroxysaiyacenol B saiyacenol C	<i>Laurencia viridis</i>	Helps in the eradication of biofilm
Proline, fucoxanthin, dimethyl sulphopropionate	<i>Fucus vesiculosus</i>	Helps in the eradication of biofilm

It was determined that the extracts obtained from approximately 225 different microalgae and cyanobacteria species showed antibiofilm activity. Among these extracts, it is known that the components showing the most antibiofilm activity are polyunsaturated fatty acids (Nag et al., 2022). It has been determined that halogenated furanones produced by seaweeds such as *D. pulchra* have an antibiofilm effect on both gram-positive and gram-negative bacterial species (Vairappan, 2003). There are studies showing that halogenated furanones also inhibit the biofilms formed by *Escherichia coli* and *P. aeruginosa*. The mechanism of inhibition occurs when furanones inhibit *P. aeruginosa* elastase B enzyme production. However, sulfur and brominated furanones were also found to inhibit biofilm formation (Eom et al., 2012).

Cepas et al. (2021) investigated the antibiofilm activities of microalgae and cyanobacteria by analytical methods such as HPLC and NMR. As a result, it has been determined that lipids obtained from cyanobacteria and microalgae show antibiofilm properties against pathogens that may cause serious infection. These lipids were thought to be an alternative source for treatments. Another study examined the antibiofilm effects of lutein from *Chlorella pyrenoidosa*, especially on resistant *P. aeruginosa*. It has been determined that even very low concentrations of lutein extract show biofilm inhibition properties. It is also thought that this may be an antiquated sensing feature. For this reason, lutein obtained from *Chlorella pyrenoidosa* can be used in the treatment of diseases such as diabetic foot ulcers, which may pose a danger due to the high biofilm formation capacity of *P. aeruginosa* (Sampathkumar et al., 2019).

3. METHODOLOGY

3.1 MICROORGANISM AND MEDIUMS

Schizochytrium sp. S31 (ATCC 20888) was used in the current investigation. Experiments on microalgal growth were conducted in a complex medium containing 15 g/l glucose and 8 g/l protease peptone, 5 g/l yeast extract, 20 g/l NaCl, and 18 g/l MOPS. On an analytical scale, all medium components were weighed, and 1 liter of distilled water was added. A homogeneous mixture was made by stirring with a magnetic stirrer. 1 ml of NaOH was added to achieve a pH of 7. The prepared broth was autoclaved for 20 minutes at 121°C.

3.2 GROWTH EXPERIMENTS

The experiments were carried out in a 250 ml flask with complex media for 48 hours at 27 °C and 200 rpm orbital shaker incubator. 500 µl of samples were taken at regular intervals under sterile conditions in a laminar flow cabinet. Using a spectrophotometer, the optical density (OD600) of 1/10 diluted samples is determined.

Spectrophotometric analysis

- a. The spectrophotometer must be prepared for use by setting and calibrating the instrument.
- b. Input a value of 600 nm for the wavelength.
- c. Resuspend the microalgal culture and take 500 µl of sample under sterile conditions. Dilute the sample 1:10 with distilled water.
- d. Read and record the absorbance of the sample at 600 nm.

3.3 DOWNSTREAM PROCESS

3.3.1 Harvesting Microalgal Culture

A microalgae culture of *Schizochytrium sp.* S31 was harvested by centrifuging at 5000 rpm for 15 minutes. After 15 minutes of centrifugation at 5000 rpm, the supernatant was discarded carefully. The cell pellet was rinsed twice with deionized water to eliminate residual medium. The lipase

enzyme, which breaks down complex lipids, was inactivated by treating the pellet in a water bath at 65°C for an hour.

3.3.2 Lyophilization

For freeze-drying the samples, a Teknosem TRS-2 stainless steel lyophilizer with a condenser temperature of -65 °C and a vacuum capacity of 2.5 m³ was employed. A harvested and cleaned microalgal cell pellet was dried at -55°C under vacuum for 70 hours.

Lyophilization Protocol:

- a. Spread the cell pellet on the lyophilizer tray and incubate it at -80 °C for 90 min.
- b. Freeze samples at -65 °C for 2 hours.
- c. Vacuum is applied under 2,5 m³ pressure. At this stage, most of the water in the sample sublimates with the heat.
- d. Air is gently introduced into the chamber to reduce pressure before opening the lid.
- e. Dried cell biomass is removed from the tray and weighted to determine CDW.

3.3.3 Cell Disruption

Different physical and chemical techniques of cell lysis were investigated to enhance the total lipid content. Two different cell lysis techniques—manual grinding and ultrasonication—were used, and the outcomes were contrasted.

3.3.3.1 Mortar and pestle

Usually, manual grinding is employed to destroy plant cells. For degradation, cell pellets were frozen at -80 °C overnight. After cooling the mortar and pestle at -20 °C for one hour, the frozen sample is pulverized using a cool mortar and pestle.

3.3.3.2 Sonication

The samples were sonicated in the extracting solvent at 50W for 15 minutes with several brief bursts (on/off every 30 seconds) by using ultrasonic homogenizer model3000. During sonication, the mixture of biomass and extraction solvent is always kept on ice to avoid overheating.

3.4 DRY BIOMASS ANALYSIS

The determination of total fat and oil (g/100g), nitrogen (protein) quantity (g/100g), energy (kcal/100g and kj/100g), carbohydrate (g/100g), moisture and/or volatile substance (g/100g), and total ash (g/100g) of the dry biomass were performed by Eurofins Izmir Food Control Laboratories, Republic of the Turkey Ministry of Agriculture and Forestry, according to Turkish Food Codex food labeling and consumer information regulation.

3.5 LIPID EXTRACTION

For the extraction of lipid from *Schizochytrium* sp. S31, solvent-based lipid extraction, which is the usual approach was applied. For lipid extraction, different solvents or solvent mixtures were used such as hexane only, ethanol only, hexane/ethanol (2:1), methanol only, and hexane/methanol (2:1). A mixture of 5 g of dried biomass and 100 ml of each solvent or solvent combination was incubated for 6 hours at room temperature and 225 rpm.

3.6 LIPID CHARACTERIZATION

3.6.1 Total Lipid Analysis

The combination of biomass and solvent was placed in a falcon tube and centrifuged to achieve a pure supernatant. Using a rotary evaporator, solvent was removed from a supernatant containing a solvent and lipid mixture. Solvent was evaporated at 95°C for 30 min. The amount of lipid was determined by subtracting the weight of the bottle before and after evaporation. The amount and proportion of lipid recovered were estimated as follows:

$$\text{Mass of lipid} = (\text{weight of the flask} + \text{extracted oil}) - (\text{weight of the flask})$$

$$\text{Lipid content (\%)} = \frac{\text{mass of lipid extracted (g)}}{\text{sample weight (g)}} \times 100$$

3.6.2 Fatty Acid Analysis

Fatty acid methyl ester (FAME) analysis was performed to analyze the samples by gas chromatography and ultimately determine the total fatty acid composition. The total lipid sample was incubated in 5 ml of 0.5N methanolic NaOH for 10 minutes at 100°C. The mixture was then mixed with 5 ml of BF₃-methanol for methylation. The extraction, evaporation, and addition of 5 ml of n-heptane, NaCl, and water resulted in the production of FAME in the top phase.

3.6.3 Gas Chromatography Analysis

FAMEs samples were transferred to the bottle containing Na₂SO₄/heptane. Gas chromatography analyses were performed by Eurofins Izmir Food Control Laboratories, Republic of Turkey Ministry of Agriculture and Forestry, according to these standards.

Conditions for the GC (Agilent Technologies 6890 N) analysis were as follows: Oven temperature was maintained at 140°C for 5 minutes then the temperature was increased to 240°C at a rate of 4°C/min. Temperatures for the detector (FID) and inlet were set at 240 to 280°C, respectively.

3.7 ANTIBACTERIAL ACTIVITY ASSAY

Overnight cultures of *Pseudomonas aeruginosa* 27853, *Escherichia coli* 35218, *Streptococcus mutans*, or *Klebsiella pneumoniae* 700603 (10⁶ CFU/ml) were incubated with *Schizochytrium* sp. S31 oil sample or uninoculated TSB medium in 96-well flat-bottom sterile plates (Nest Scientific, Maryland, USA) at 37 °C for 8 or 24 hours without shaking. A multiwell plate reader (Multiskan FC, USA) was used to measure the rate of bacterial growth at a wavelength of 600 nm. The optical density of the uninoculated bacterial growth medium, and medium containing a *Schizochytrium* sp. S31 oil sample at 600 nm were used as blank. For the plate assays, *P. aeruginosa*, *Escherichia coli* 35218, *Streptococcus mutans*, or *K. pneumonia* was grown in *Schizochytrium* sp. S31 oil samples (or in control growth medium; TSB or medium containing *Schizochytrium* sp. S31 oil sample) in 96-well plates then, sample were serially diluted by using TSB medium. 3 µl of the diluted samples were taken and pipetted into agar plates and incubated at 37°C overnight. The spots were visualized using the G-BOX imaging system.

3.8 BIOFILM FORMATION ASSAY

The biofilm formed on the 96-well plate was evaluated with the crystal violet (CV) staining protocol, as described previously (Merritt et al. 2011; Tunçer & Karacam, 2020). *Schizochytrium* sp. S31 oil sample was inoculated with *P. aeruginosa* and *Escherichia coli* 35218 (10^6 CFU/ml) in 96-well f-bottom polystyrene plates (Nest Scientific). The plate samples were incubated for 24 hours at 37°C without shaking, then all the medium in the well was removed and washed three times to get rid of any remaining planktonic bacteria. For the fixation process, 150 µl of methanol were added to each well, and incubated at room temperature for 15 minutes. After methanol was removed, the plates were dried by air. The plates were incubated at room temperature for 15 minutes after being stained with 130 µl of a 0.1% crystal violet. The stain were removed and the plates were washed three times to remove any CV residues. 130 µl of 30% acetic acid was added to each well of plates for solubilization of the CV. Optical density at 550 nm was measured using a multi-well plate reader. The uninoculated culture was used as a control.

4. RESULTS

4.1 GROWTH EXPERIMENTS

Schizochytrium sp. S31, a stock culture, was stored at -80 °C with a mixture of cryoprotectant consisting of 5% DMSO and 5% glycerol. The viability checks of the stocks were carried out at regular intervals. Under sterile conditions, the sample taken from the stock with a loop was inoculated on a complex agar medium and incubated at 25 °C for 72 hours. Colonies were counted after the incubation. In cases of small numbers of colonies, stock cultures were re-prepared from fresh cultures. For microalgae growth curve analysis, a single colony was selected and inoculated into 50 ml of sterile complex medium in a 250 ml flask. The culture was incubated for 48 hours at 25°C at 225rpm. The growth curve of *Schizochytrium* sp. S31 was obtained by measuring the optical density of the samples taken from the culture at regular intervals at OD₆₀₀ nm using a spectrophotometer. Figure 4.1 shows the growth curve of *Schizochytrium* sp. S31 in a complex medium.




Figure 4.1: Growth curve of *Schizochytrium* sp. S31

Figure 4.1 shows that *Schizochytrium* sp. S31 grows logarithmically between 4-24 hours before entering the stationary phase at 24 hours due to the decrease in the amount of nutrients.

4.2 DRY BIOMASS CALCULATION AND ANALYSIS

Microalgae cell culture was harvested by centrifuging at 5000 rpm for 15 minutes, and the supernatant was discarded. The cell pellet was washed twice with sterile deionized water. The pellet was placed in a 65°C water bath for one hour to inactivate the lipase enzyme. The harvested and washed cell pellet was lyophilized at -55°C under vacuum for 70 hours by following the freeze-dry protocol. The lyophilized biomass was scraped off the tray and powdered using a simple homogenizer. Cell dry weight was calculated as the amount of weighed biomass per liter (g/L) (Table 4.1).

Table 4.1: Chemical characteristics of the of *Schizochytrium* sp. S31 dry biomass

Analyses	Result
Cell Dry Weight (g/L)	57.1
Determination of Total Fat and Oil) (g/100g)	5.31
Determination of Nitrogen (Protein) Quantity) (g/100g(Nx6,25)	44.62
Determination of Energy (kcal / 100g)	374.95
Determination of Energy (kj/100g)	1568.79
(Determination of Moisture and / or Volatile Substance, Drying loss, Moisture content) (g/100g)	7.19
(Total ash) (g/100g)	5.71

Table 4.1 shows the amount of total fat and oil (g/100g), nitrogen (protein) quantity (g/100g), energy (kcal/100g and kj/100g), carbohydrate (g/100g), moisture and / or volatile substance (g/100g) and total ash (g/100g) of the dry biomass determined by Eurofins Izmir Food Control Laboratories, Republic of Turkey Ministry of Agriculture and Forestry. The analysis report is placed in Appendix A.1. According to the results, the weight of dry biomass with a moisture content of 7.19% was calculated as 57.1 g/l. Dry biomass is composed of 44.62% of protein and 5.31% of total fat.

Total amino acid and vitamin amounts of dry biomass were shown in Tables 4.2 and 4.3, respectively. Analysis report is placed in Appendix A.2.

Table 4.2: Total amino acid content of the dry biomass

Amino Acid	g/100g protein
Aspartic acid	10.5 ± 0.09
Threonine	5.24 ± 0.12
Serine	5.08 ± 0.07
Glutamic acid	10.7 ± 0.03
Proline	6.45 ± 0.26
Glycine	5.13 ± 0.07
Alanine	8.44 ± 0.19
Valine	6.44 ± 0.12
Methionine	1.52 ± 0.13
Isoleucine	5.01 ± 0.14
Leucine	6.84 ± 0.17
Tyrosine	5.21 ± 0.21
Phenylalanine	4.26 ± 0.04
Histidine	5.02 ± 0.10
Lysine	5.63 ± 0.08
Arginine	8.25 ± 0.26

According to the amino acid content of the proteins in dry biomass, the amino acids with the highest ratio are responsible for aspartic acid and glutamic acid, with the ratios of 10.5g/ 100g and 10.7g/ 100g, respectively.

Table 4.3: Total vitamin content of the dry biomass

Vitamin	Concentration
Iron, mg/kg	869.51 ± 38.96
Niacin, mg/kg	245 ± 16.58
Potassium, mg/kg	1360 ± 241.74
Calcium, mg/kg	100.08 ± 9.62
Lutein, mg/kg	368.65 ± 56.72
Vitamin A, IU/kg	216.41 ± 72.13
Vitamin B 1, mg/kg	4.48 ± 0.12
Vitamin B 2, mg/kg	14.26 ± 3.04
Vitamin B 6, mg/kg	9.62 ± 2.84
Vitamin C, mg/kg	74.16 ± 1.85
Vitamin B 12, µg/kg	725.17 ± 156.20

Potassium (1360 mg/kg), iron (869.51 mg/kg), vitamin B12 (725.17 mg/kg), lutein (368.65 mg/kg), niacin (245 mg/kg), vitamin A (216.41 mg/kg), and calcium (100.08 mg/kg) were found to have the highest quantities of vitamins and elements in dry biomass.

4.3 COMPARISON OF SEVERAL CELL LYSIS AND LIPID EXTRACTION TECHNIQUES

Cell lysis and lipid extraction are operations that are a part of the downstream process. For cell lysis, two distinct techniques—mortar and pestle and ultrasonication—were evaluated, and the results were then compared. Five different solvent mixtures, hexane only, ethanol only, and hexane/ethanol (2:1), methanol only, hexane/methanol (2:1), were examined for lipid extraction.

When hexane/methanol (2:1) was used among five different solvents, the best result was obtained with a 34% lipid ratio. The outcomes were averaged after each experiment was carried out three

times. Error bars show the standard deviation. Comparative results of the cell lysis and lipids were shown in Figure 4.2.




Figure 4.2: Comparative results of different cell lysis and lipid extraction methods

4.4 GAS CHROMATOGRAPHY ANALYSIS

Gas chromatography analyses were performed by Eurofins Izmir Food Control Laboratories, Republic of Turkey Ministry of Agriculture and Forestry, according to standards. The analysis report is placed in Appendix A.3. The fatty acid profile results obtained from GCMS analysis were shown in Table 4.4.

Table 4.4: Fatty acid composition of *Schizochytrium* sp. S31

Analyses	Result
Arachidic Acid (C20:0)	0.41
Arachidonic Acid (C20:4)	1.07
Behenic acid(C22:0)	0.25
Eicosenoic Acid (C20: 1)	1.15
Eicosatrienoic Acid (C20: 3 (n3)	0.61
Eicosatrienoic Acid (C20: 3(n6)	0
Eicosadienoic Acid (C20:2)	0.17
Docosahexaenoic Acid (DHA) (C22: 6)	10.47
Eicosapentaenoic Acid (C22:5)	1.84
Eicosapentaenoic Acid (EPA) (C20:5)	16.11
Gamma Linolenic Acid (C18:3 (n6)	0
Heneicosanoic Acid (C21:0)	2.93
Heptadecanoic Acid (C17:0)	0.76
Heptadecenoic Acid (C17:1)	0.05
Capric Acid (C10:0)	0
Caprylic Acid (C8:0)	0
Lauric acid (C12:0)	0.17
Lignoceric Acid (C24:0)	0.78
Erucic Acid (C22:1)	0.11
Linoleic Acid (C18:2)	1.49
Myristic Acid (C14:0)	7.92
Myristoleic Acid (C14:1)	0.23
Nervonic Acid (C24:1)	0.33
Oleic Acid (C18:1)	16.8
Palmitic Acid(C16:0)	14.38
Penta-Decanoic Acid (C15:0)	0.49
Palmitoleic Acid (C16:1)	10.11
Stearic Acid (C18:0) Total	2.84
Omega-3 + Omega-6 (Fatty Acids)	32.15
Total Omega-3 Fatty Acids	27.42
Total Omega-6 Fatty Acids	4.73
Trans Linolenic Acid (tC18:3)	0.13
Trans-9-Octadecenoic Acid (Trans Oleic Acid) (tC18:1)	0.15
Trans-9,12- Octadecenoic Acid (Trans Linoleic Acid) (tC18:2)	0.06
Tricosanoic Acid (C23:0)	0.75
Unidentified Fatty Acids	6.38

Omega-3 + Omega-6 (Fatty Acids) lipid ratio had the highest rate at 32.15%, according to the results. The ratio of total omega-3 fatty acids, which is the second-highest lipid ratio, was found to be 27.42%. The ratios of Oleic acid (C18:1), Eicosapentaenoic Acid (EPA) (C20:5), Palmitic Acid (C16:0), Docosahexaenoic acid (DHA) (C22:6), which are the fatty acids with the highest values in the total fatty acid profile, were obtained as 16.8%, 16.11%, 14.38%, and 10.47%, respectively. Palmitoleic acid (C16:1) and Myristic Acid (C14:0) fatty acids are intermediate-level fatty acids obtained at 10.11% and 7.92%, respectively. Heneicosanoic Acid (C21:0), Stearic Acid (C18:0), Eicosapentaenoic Acid (C22:5), and Linoleic Acid (C18:2) are low-level fatty acids obtained at 2.93%, 2.84%, 1.84%, and 1.49%, respectively. Apart from these, Arachidic Acid (C20:0), Arachidonic Acid (C20:4), Behenic acid (C22:0), Eicosenoic Acid (C20:1), Eicosatrienoic Acid (C20: 3), Eicosadienoic Acid (C20:2), Gamma Linolenic Acid (C18:3), Heptadecanoic Acid (C17:0), Heptadecenoic Acid (C17:1), Capric Acid (C10:0), Caprylic Acid (C8:0), Lauric acid (C12:0), Lignoceric Acid (C24:0), Erucic Acid (C22:1), Myristoleic Acid (C14:1), Nervonic Acid (C24:1), Penta-Decanoic Acid (C15:0), Trans Linolenic Acid (tC18:3), Trans-9-Octadecenoic Acid (Trans Oleic Acid) (tC18:1), Trans-9,12- Octadecenoic Acid (Trans Linoleic Acid) (tC18:2) and Tricosanoic Acid (C23:0) are in the class of fatty acids that are almost absent in their fatty acid composition (Table 4.4).

4.5 ANTIBACTERIAL ACTIVITY

Schizochytrium sp. S31 oil sample was incubated with 10^6 CFU/ml of three different gram-negative pathogen bacteria, *Escherichia coli* 35218, *Pseudomonas aeruginosa* 27853, *Klebsiella pneumoniae* 700603, as well as the Gram-positive pathogen *Streptococcus mutans*, at 37 °C for 8 or 24 hours without shaking. Uninoculated TSB medium was used as a negative control. The bacterial growth was determined by using a multi-well plate reader at OD₆₀₀. For spot plate assays, undiluted and serially diluted pathogenic bacterial samples were incubated separately for 8 and 24 hours with *Schizochytrium* sp. oil samples at various rates. The spots were captured on camera using the G: Box imaging technology. The experiments were repeated three times, and the results from each trial are shown as SEM. The t test was used for the analysis of the results. The antibacterial effect of a *Schizochytrium* sp. S31 oil sample on *Pseudomonas aeruginosa* 27853, *Escherichia coli* 35218, *Streptococcus mutans*, and *Klebsiella pneumoniae* 700603 were shown in Figure 4.3, Figure 4.4, Figure 4.5, and Figure 4.6, respectively.




Figure 4.3: Investigation of antibacterial effect of *Schizochytrium* sp. S31 oil sample on *P. aeruginosa* bacteria. *P. aeruginosa* was incubated with oil or with control medium, for **a.** 8h and **b.** 24h. The bacterial growth was determined at OD600. Spot plate assays for **c.** 8h and **d.** 24h were used to visualizing cell growth. Then incubated at 37°C. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).




Figure 4.4: Investigation of antibacterial effect of *Schizochytrium* sp. S31 oil sample on *Escherichia coli* 35218. *E. coli* was incubated with oil or with control medium, for a. 8h and b. 24h. The bacterial growth was determined at OD600. Spot plate assays for c. 8h and d. 24h were used to visualizing cell growth. Then incubated at 37. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$)




Figure 4.5: Investigation of antibacterial effect of *Schizochytrium* sp. S31 oil sample on *Streptococcus mutans*. *S. mutans* was incubated with oil or with control medium, for a. 8h and b. 24h. The bacterial growth was determined at OD600. Spot plate assays for c. 8h and d. 24h were used to visualizing cell growth. Then incubated at 37. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).




Figure 2.6: Investigation of antibacterial effect of *Schizochytrium* sp. S31 oil sample on *Klebsiella pneumoniae*. *K. pneumoniae* was incubated with oil or with control medium, for **a**. 8h and **b**. 24h. The bacterial growth was determined at OD600. Spot plate assays for **c**. 8h and **d**. 24h were used to visualizing cell growth. Then incubated at 37. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).

According to results, the *Schizochytrium* sp. S31 oil sample was found to have antibacterial activity on pathogenic bacteria of *Pseudomonas aeruginosa*, *Escherichia coli* 35218, *Klebsiella pneumonia*, and *Streptococcus mutans*.

Antibacterial activity study on *Pseudomonas aeruginosa* showed that cell viability decreased by 1.3, 2.3, 4.0, and 7.7 times, when the bacterial culture was incubated for 8 hours with 10%, 20%,

40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively. Cell viability was determined to decrease by 1.9 and 1.7 times, at 40% and 60% concentration of *Schizochytrium* sp. S31 oil respectively, when the bacterial culture was treated with for 24 hours.

Antibacterial activity study on *Escherichia coli* 35218 showed that cell viability decreased by 1.1, 1.4, 2.3, and 13.1 times, when the bacterial culture was incubated for 8 hours with 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively. When the bacterial culture was incubated with *Schizochytrium* sp. S31 oil at 40% and 60% concentration for 24 hours, cell viability was observed to decrease by 1.9 and 2.4 times, respectively.

Antibacterial activity study on *Klebsiella pneumonia* showed that cell viability decreased by 1.4, 3.7, 6.7, and 72.7 times, when the bacterial culture was incubated for 8 hours with 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively. Cell viability was shown to decrease by 1.4, 1.7, 14.3, and 9.7 times, at 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil when the bacterial culture was treated with for 24 hours respectively.

Antibacterial activity study on *Streptococcus mutans* showed that cell viability decreased by 1.1, 1.8, 2.9, and 6.4 times, when the bacterial culture was incubated for 8 hours with 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively. When the bacterial culture was incubated for 24 hours, cell viability was observed to decrease by 1.3, 2.5, and 15.8 times, at 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively.

4.6 ANTIBIOFILM ACTIVITY

Schizochytrium sp. S31 oil sample was incubated with 10^6 CFU/ml of two different gram-negative pathogen bacteria, *Escherichia coli* 35218 and *Pseudomonas aeruginosa* 27853 in 96-well flat-bottom polystyrene cell culture plates at 37 °C for 8 or 24 hours without shaking. The samples were washed, dried and then stained with crystal violet. After staining, plates were washed three times to remove any residual CV. Optical density at 550nm was measured using a multi-well plate reader. The uninoculated culture was used as a control. Effect of *Schizochytrium* sp. S31 oil sample on biofilm formation in *P. aeruginosa* and *Escherichia coli* 35218, were shown in Figure 4.6 and Figure 4.7 respectively. According to results, surface adhesion ability and biofilm forming capacity of *P. aeruginosa*, a widely used model bacterium in biofilm studies, were severely inhibited in the

presence of *Schizochytrium* sp. S31 oil sample. The same results were observed with the known biofilm-forming gram-negative bacteria, *Escherichia coli* 35218.




Figure 4.7: Effect of *Schizochytrium* sp. S31 oil sample on biofilm formation in *P. aeruginosa*. Biofilm development in *P. aeruginosa*, which was incubated with the oil used in the specified amounts for a. 8h and b. 24h, c. 8h d. 24h was investigated by CV staining. Quantitative results are given in percent (%) compared to *P. aeruginosa* grown in control medium. Well staining are representative experimental results. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).




Figure 4.8: Effect of *Schizochytrium* sp. S31 oil sample on biofilm formation in *Escherichia coli* 35218. Biofilm development in *E. coli*, which was incubated with the oil used in the specified amounts for a. 8h and b. 24h, c. 8h d. 24h was investigated by CV staining. Quantitative results are given in percent (%) compared to *Escherichia coli* grown in control medium. Well staining are representative experimental results. Experiments were repeated three times and results from each trial are shown as $\pm$ SEM. The t test was used for the analysis of the results (ns:no significant** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$).

The antibiofilm activity investigation on *P. aeruginosa* revealed that biofilm formation was decreased by 22.3%, 76.3%, 88.3%, and 91% when the bacterial culture was treated for 8 hours and by 46.5%, 52.7%, 78.8%, and 96% when it was treated for 24 hours with 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively.

According to the results of the antibiofilm activity study on *Escherichia coli*, biofilm formation decreased by 42.5%, 71.3%, 84.9%, and 94.7%, when the bacterial culture was treated for 8 hours and by 25.4%, 51%, 78.7%, and 89.5%, when it was treated for 24 hours with 10%, 20%, 40%, and 60% concentration of *Schizochytrium* sp. S31 oil respectively.



5. DISCUSSION AND CONCLUSION

Algae are harmless and effective sources of bioactive molecules with a wide range of uses. Algae themselves or their extracts have the potential to be used in many different areas. Food additive, animal feed, aquaculture, biofertilizer, drug precursor, anti-inflammatory, anti-allergic, and supplement drugs, as well as antimicrobial and anti-biofilm products, waste treatment, bioenergy, and CO₂ emission (thus a potential global warming prevention agent) are a few examples of applications. (Concas et al., 2013; Ji et al., 2014; Axelsson et al., 2012). Algae can be used as a bio-based compound or as biodiesel from raw materials or refined products (Bahadar & Khan, 2013). Also, it has been demonstrated that chemical compounds derived from algae, such as lipids, proteins, enzymes, vitamins, and antioxidant chemicals, are beneficial for human health (Parsaeimehr et al., 2015). Biological compounds such as EPA, DHA, astaxanthin, and protein pigments produced by algae are widely used in the health, pharmaceutical, and food sectors. With all these features and the widespread use of algae, many private companies from all over the world have started to invest much more in algae-based bioproducts. UAS facilities and research programs provide potential for sustainable energy, sustainable fuel, food, feed, and many other emerging algae-based businesses. Solazyme Roquette Company, located in San Francisco, America produces microalgae and products. This company has recently launched a product called Almagine HL. This product is a natural, environmentally friendly, nutritious, rich food containing micronutrients including carbohydrates (simple sugars and polysaccharides), protein, lipids, fiber and lutein. Commercialization of algae products has accelerated in recent years, and many new products will be produced in the coming years. Moreover, numerous species of algae and microalgae have been studied recently for their antibacterial, antifungal, antiviral, and antibiofilm properties, and remarkable results have been obtained (Desbois et al., 2008; Smith et al., 2010; Veetil et al., 2022; Santoyo et al., 2009; Abedin & Taha, 2008; Katircioğlu et al., 2006). Due to the increasing need and the insufficient effect of existing antimicrobial agents, microalgae extract can also be used in treatment due to its antimicrobial activities.

In this study, the antimicrobial and antibiofilm activity of heterotrophic microalgae *Schizochytrium* sp. S31 microalgae oil were investigated. For this purpose, *Schizochytrium* sp. S31 was first cultivated heterotrophically under optimum conditions. Microalgae can grow rapidly under controlled conditions that provides great advantages. The level of some macromolecules in the

biomass can also be enhanced when the growth conditions are changed as desired. For example, it is known that the polar lipid content of microalgae increases under the low temperature and nitrogen-limiting environment (Vardar et al., 2016). Microalgae cell culture was then harvested and dried by lyophilization. Cell dry weight (CDW) was calculated as 57.1 g/l. In the downstream processing stage, various cell lysis and lipid extraction procedures were examined. Five different solvent mixtures—hexane, hexane/ethanol (2:1), ethanol, methanol, and hexane/methanol (2:1)—as well as two different cell disruption techniques—mortar and pestle and ultrasonication—were examined and compared. The total lipid yields were increased up to 34% with a clear appearance using hexane/methanol (2:1) together with ultrasonication. The extraction of high yield lipid from *Schizochytrium* sp. microalgae has previously been studied using a variety of cell lysis and lipid extraction procedures. Vardar-Yel et al. (2017) examined various cell lysis (French press, ultrasonication, CTAB, SDS, alcalase) and lipid extraction (hexane, isopropanol, ethanol, and supercritical CO₂) methods on *Schizochytrium* sp. and discovered that when hexane solvent with sonication and/or French press was used, a high content of lipid with a high TAG content was obtained. In another study, acid treatment, enzyme digestion, osmotic shock, and homogenizer methods were combined with the Bligh and Dyer or Soxhlet methods to obtain high lipid content from *Schizochytrium* sp. It was stated that the most efficient result was obtained by increasing the lipid ratio above 21% and the %DHA to 19.25% as a result of the application of the Bligh and Dyer method together with the enzyme application (Hac Isa et al., 2022). Byreddy et al., (2015) carried out a similar study with *Schizochytrium* sp. S31 and *Thraustochytrium* sp. AMCQS5-5 strains using various solvent combinations. The author also tried different cell lysis methods such as grinding with liquid nitrogen, a water bath, osmotic shock, bead vortexing, and sonication. They obtained the highest lipid ration (48.7%) by using an osmotic shock method together with chloroform:methanol (2:1). Gas chromatography was used to assess the fatty acid content following lipid extraction. According to the GCMS analysis results, 10.4% DHA and 16.11% EPA fatty acids were obtained. Apart from these, other fatty acids with high levels were determined as 16.8% oleic acid, 14.8% palmitic acid and 10.11% palmitoleic acid. Total Omega-3 Fatty Acids and Total Omega-3 + Omega-6 Fatty Acids ratio were calculated as 27.42% and 32.15% respectively. *Schizochytrium* sp. S31 is once again confirmed to be rich in omega-3 fatty acids. Numerous other research in the literature demonstrate that *Schizochytrium* sp. contains a high concentration of omega-3 fatty acids. (Hakim, 2012; Ward et al., 2005; Byreddy, 2016).

P. aeruginosa is a pathogenic bacterium that can form biofilms on biotic and abiotic surfaces and is used as a model for biofilm formation (Pericolini et al., 2018). *P. aeruginosa*, an opportunistic pathogenic bacterium, can cause serious infections, including pneumonia, urinary tract infections, otitis externa, otitis media, soft tissue infections (keratitis), and bacteremia (Lyczak et al., 2000). Infections caused by *P. aeruginosa* often lead to high morbidity and mortality due to resistance to antibiotics and antimicrobial treatments. *P. aeruginosa* infections are generally characterized by eliciting an intense neutrophilic response (Gellatly & Hancock, 2013). *K. pneumonia* also causes many human infections. These include various infections such as bacteremia, pneumonia, urinary tract infections, meningitis, bloodstream infection, and intra-abdominal infection (Caneiras et al., 2019). These bacteria can be commonly found in raw fish, meat, milk, and dairy products, as well as in street foods (Zhang et al., 2018). *Escherichia coli* (*E. coli*) is a pathogenic bacterium known to cause gastrointestinal diseases such as diarrhea in humans and animals. *E. coli* has the capacity to easily form biofilms on the surface of living or non-living carriers, which can lead to cross-contamination in foods (Zhou et al., 2022). *Streptococcus mutans* (*S. mutans*) are gram-positive bacteria found in the human mouth and more specifically on tooth surfaces. *S. mutans* are able to produce large quantities of glucans and acids, exceeding their saliva buffering capacity, giving bacteria the advantage to outcompete non-cariogenic common species in low pH environments (Metwalli et al., 2013).

In this study, the antibacterial effects of *Schizochytrium* sp. S31 oil samples were investigated on Gram-negative pathogens, which cause many different human infections, as well as the Gram-positive pathogen *Streptococcus mutans*. *Schizochytrium* sp. S31 oil sample was found to have antibacterial activity on pathogenic bacteria including *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, and *Escherichia coli* 35218. It can be concluded that the best results in terms of antibacterial activity were obtained under the condition of 60% concentration of *Schizochytrium* sp. S31 oil for 8 hours for *Pseudomonas aeruginosa*, *Escherichia coli* 35218 and *Klebsiella pneumonia* with a decrease of 7.7%, 13.1% and 72.7% in cell viability respectively. In the study with *Streptococcus mutans*, the highest antibacterial activity was obtained with a decrease of 15.8% in cell viability under 60% concentration of *Schizochytrium* sp oil for 24 hours. Although it was clear from all of the results that the oil sample had an antibacterial effect on all the tested pathogenic organisms, the most striking outcome came from the treatment of the 60% concentration *Schizochytrium* sp. S31 oil sample with *Klebsiella pneumonia* for 8 hours.

Antibiofilm activity of the *Schizochytrium* sp. S31 oil on *Pseudomonas aeruginosa*, and *Escherichia coli* 35218 bacteria were also investigated. Both pathogens are also known for their biofilm-forming potential. It is possible to conclude that the conditions for the optimum antibiofilm activity were 60% *Schizochytrium* sp. S31 oil concentration for 24 hours for *Pseudomonas aeruginosa* and 8 hours for *Escherichia coli* 35218 with a decrease of 96% and 94.7% in terms of biofilm formation, respectively.

As a conclusion, the *Schizochytrium* sp. S31 oil sample was found to have anti-pathogenic activity on pathogenic bacteria of *Pseudomonas aeruginosa*, *Escherichia coli* 35218, *Klebsiella pneumonia* and *Streptococcus mutans*. In addition, the surface adhesion ability and biofilm formation capability of *P. aeruginosa*, a frequently used model bacteria in biofilm research, were investigated (Ryder et al., 2007), and were dramatically inhibited in the presence of a *Schizochytrium* sp. S31 oil sample. The same results were obtained with the gram-negative bacteria *Escherichia coli* 35218 strain, which forms biofilms (Mariscal et al., 2007). Although the active compound in the *Schizochytrium* sp. S31 oil sample with antibacterial and antibiotic capabilities is not yet known, it is assumed to be DHA and EPA, which have the highest ratio in fatty acid composition. In future studies, it can be determined which active substance with antibacterial and antibiofilm properties in *Schizochytrium* sp. S31 oil sample. This study highlights that *Schizochytrium* sp. S31 oil can potentially be a natural antibacterial and antibiofilm agent that could be applied in the pharmaceutical and food industries.

REFERENCES

- Abd El-Baky, H. H., & El-Baroty, G. S. (2013). The potential use of microalgal carotenoids as dietary supplements and natural preservative ingredients. *Journal of Aquatic Food Product Technology*, 22(4), 392–406.
- Abedin, R. M., & Taha, H. M. (2008). Antibacterial and antifungal activity of cyanobacteria and green microalgae. Evaluation of medium components by Plackett-Burman design for antimicrobial activity of *Spirulina platensis*. *Global Journal of Biotechnology and Biochemistry*, 3(1), 22-31.
- Aguilar-Rosas, L. E., Núñez-Cebrero, F., & Aguilar-Rosas, C. (2013). Introduced marine macroalgae in the Port of Ensenada, Baja California, Mexico: biological contamination. *Procedia Environmental Sciences*, 18, 836-843.
- Amaro, H. M., Guedes, A. C., & Malcata, F. X. (2011). Antimicrobial activities of microalgae: an invited review. *Science against microbial pathogens: communicating current research and technological advances*, 3, 1272-1284.
- Ambati, R. R., Siew Moi, P., Ravi, S., & Aswathanarayana, R. G. (2014). Astaxanthin: Sources, extraction, stability, biological activities and its commercial applications—A review. *Marine drugs*, 12(1), 128-152.
- Anaissie, E., & Bodey, G. P. (1989). Nosocomial fungal infections: old problems and new challenges. *Infectious disease clinics of North America*, 3(4), 867-882.
- Apt, K. E., & Behrens, P. W. (1999). Commercial developments in microalgal biotechnology. *Journal of phycology*, 35(2), 215-226.
- Axelsson, L., Franzén, M., Ostwald, M., Berndes, G., Lakshmi, G., & Ravindranath, N. H. (2012). *Jatropha* cultivation in southern India: assessing farmers' experiences. *Biofuels, Bioproducts and Biorefining*, 6(3), 246-256.
- Baweja, P., Sahoo, D. (2015). Classification of Algae. D. Sahoo, J. Seckbach (Ed.), *The Algae World* (31-55. ss.). Dordrecht: Springer. https://doi.org/10.1007/978-94-017-7321-8_2

- Bahadar, A., & Khan, M. B. (2013). Progress in energy from microalgae: a review. *Renewable and Sustainable Energy Reviews*, 27, 128-148.
- Batinić, D., & Robey, F. A. (1992). The V3 region of the envelope glycoprotein of human immunodeficiency virus type 1 binds sulfated polysaccharides and CD4-derived synthetic peptides. *Journal of Biological Chemistry*, 267(10), 6664-6671.
- Becker, E. W. (2007). Micro-algae as a source of protein. *Biotechnology Advances*, 25(2), 207–210.
- Benkendorff, K., Davis, A. R., Rogers, C. N., & Bremner, J. B. (2005). Free fatty acids and sterols in the benthic spawn of aquatic molluscs, and their associated antimicrobial properties. *Journal of Experimental Marine Biology and Ecology*, 316(1), 29–44.
- Bhagavathy, S., Sumathi, P., & Bell, I. J. S. (2011). Green algae *Chlorococcum humicola*-a new source of bioactive compounds with antimicrobial activity. *Asian Pacific Journal of Tropical Biomedicine*, 1(1), 1-7.
- Blockx, J., Verfaillie, A., Thielemans, W., & Muylaert, K. (2018). Unravelling the mechanism of chitosan-driven flocculation of microalgae in seawater as a function of pH. *ACS Sustainable Chemistry & Engineering*, 6(9), 11273-11279.
- Borowitzka, M. A. (1998). Algae as food. B.J. Wood (Ed.) *Microbiology of Fermented Foods* (585–602. ss.). London: Thomson Science. https://doi.org/10.1007/978-1-4613-0309-1_18.
- Borowitzka, M. A. (2013). High-value products from microalgae-their development and commercialisation. *Journal of Applied Phycology*, 25(3), 743–756.
- Borowitzka, M. A., & Larkum, A. W. D. (1987). Calcification in algae: mechanisms and the role of metabolism. *Critical Reviews in Plant Sciences*, 6(1), 1–45.
- Borowitzka, M. A., & Vonshak, A. (2017). Scaling up microalgal cultures to commercial scale. *European Journal of Phycology*, 52(4), 407–418.
- Bourgougnon, N., & Stiger-Pouvreau, V. (2011). Chemodiversity and bioactivity within red and brown macroalgae along the French coasts, metropole and overseas departements and territories. *Se-*

- Kwon Kim (Ed.), Handbook of marine macroalgae: biotechnology and applied phycology (58-105. ss.). New Jersey: John Wiley & Sons, Inc. <https://doi.org/10.1002/9781119977087.ch4>
- Byreddy, A. R., Gupta, A., Barrow, C. J., & Puri, M. (2015). Comparison of cell disruption methods for improving lipid extraction from thraustochytrid strains. *Marine drugs*, 13(8), 5111-5127.
- Byreddy, A. R. (2016). Thraustochytrids as an alternative source of omega-3 fatty acids, carotenoids and enzymes. *Lipid Technology*, 28(3-4), 68-70.
- Calder, P. C. (2013). Omega-3 polyunsaturated fatty acids and inflammatory processes: Nutrition or pharmacology? *British Journal of Clinical Pharmacology*, 75(3), 645–662.
- Caneiras, C., Lito, L., Melo-Cristino, J., & Duarte, A. (2019). Community-and hospital-acquired *Klebsiella pneumoniae* urinary tract infections in Portugal: virulence and antibiotic resistance. *Microorganisms*, 7(5), 138.
- Cavalier-Smith, T. (1999). Principles of protein and lipid targeting in secondary symbiogenesis: Euglenoid, dinoflagellate, and sporozoan plastid origins and the eukaryote family tree. *Journal of Eukaryotic Microbiology*, 46(4), 347–366.
- Cardozo, K. H., Guaratini, T., Barros, M. P., Falcão, V. R., Tonon, A. P., Lopes, N. P., ... & Pinto, E. (2007). Metabolites from algae with economical impact. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 146(1-2), 60-78.
- Cepas, V., Gutiérrez-Del-Río, I., López, Y., Redondo-Blanco, S., Gabasa, Y., Iglesias, M. J., ... & Soto, S. M. (2021). Microalgae and cyanobacteria strains as producers of lipids with antibacterial and antibiofilm activity. *Marine Drugs*, 19(12), 675.
- Certik, M., & Shimizu, S. (1999). Biosynthesis and regulation of microbial polyunsaturated fatty acid production. *Journal of Bioscience and Bioengineering*, 87(1), 1–14.
- Chanda, W., Joseph, T. P., Guo, X. F., Wang, W. D., Liu, M., Vuai, M. S., Padhiar, A. A., & Zhong, M. T. (2018). Effectiveness of omega-3 polyunsaturated fatty acids against microbial pathogens. *Journal of Zhejiang University: Science B*, 19(4), 253–262.

- Cheng, D., Li, D., Yuan, Y., Zhou, L., Li, X., Wu, T., Wang, L., Zhao, Q., Wei, W., & Sun, Y. (2017). Improving carbohydrate and starch accumulation in *Chlorella* sp. AE10 by a novel two-stage process with cell dilution. *Biotechnology for Biofuels*, 10(1), 1–14.
- Chi, Z., Liu, Y., Frear, C., & Chen, S. (2009). Study of a two-stage growth of DHA-producing marine algae *Schizochytrium limacinum* SR21 with shifting dissolved oxygen level. *Applied Microbiology and Biotechnology*, 81(6), 1141–1148.
- Corrêa, D. de O., Duarte, M. E. R., & Nosedá, M. D. (2019). Biomass production and harvesting of *Desmodesmus subspicatus* cultivated in flat plate photobioreactor using chitosan as flocculant agent. *Journal of Applied Phycology*, 31(2), 857–866.
- Concas, A., Lutz, G. A., Locci, A. M., & Cao, G. (2013). *Nannochloris eucaryotum* growth: Kinetic analysis and use of 100% CO₂. *Advances in Environmental Research*, 2(1), 19–33.
- Dai, L., Tan, L., Jin, X., Wu, H., Wu, H., Li, T., & Xiang, W. (2020). Evaluating the potential of carbohydrate-rich microalga *Rhodospira* sp. SCSIO-45730 as a feedstock for biofuel and β -glucans using strategies of phosphate optimization and low-cost harvest. *Journal of Applied Phycology*, 32(5), 3051–3061.
- Damatac, A. M., & Cao, E. P. (2022). Identification and diversity assessment of cyanobacterial communities from some mine tailing sites in Benguet Province, Philippines using isolation-dependent and isolation-independent methods. *Environment, Development and Sustainability*, 24(1), 1166–1187.
- De Oliveira, M. A. C. L., Monteiro, M. P. C., Robbs, P. G., & Leite, S. G. F. (1999). Growth and chemical composition of *Spirulina maxima* and *Spirulina platensis* biomass at different temperatures. *Aquaculture International*, 7(4), 261–275.
- Desbois, A. P., Lebl, T., Yan, L., & Smith, V. J. (2008). Isolation and structural characterisation of two antibacterial free fatty acids from the marine diatom, *Phaeodactylum tricornutum*. *Applied microbiology and biotechnology*, 81(4), 755–764.
- Desbois, A. P., & Smith, V. J. (2010). Antibacterial free fatty acids: Activities, mechanisms of action and biotechnological potential. *Applied Microbiology and Biotechnology*, 85(6), 1629–1642.

- Desbois, A. P., Mearns-Spragg, A., & Smith, V. J. (2009). A fatty acid from the diatom *Phaeodactylum tricornutum* is antibacterial against diverse bacteria including multi-resistant *Staphylococcus aureus* (MRSA). *Marine Biotechnology*, 11(1), 45–52.
- Dunahay, T. G., Jarvis, E. E., Dais, S. S., & Roessler, P. G. (1996). Manipulation of microalgal lipid production using genetic engineering. *Applied Biochemistry and Biotechnology - Part A Enzyme Engineering and Biotechnology*, 57, 223–231.
- Eom, S. H., Kim, Y. M., & Kim, S. K. (2012). Antimicrobial effect of phlorotannins from marine brown algae. *Food and Chemical Toxicology*, 50(9), 3251–3255.
- Feldlaufer, M. F., Knox, D. A., Lusby, W. R., & Shimanuki, H. (1993). Antimicrobial activity of fatty acids against *Bacillus larvae*, the causative agent of American foulbrood disease. *Apidologie*, 24(2), 95–99.
- Ehresmann, D. W., Deig, E. F., & Hatch, M. T. (1979). Anti-viral properties of algal polysaccharides and related compounds. T. Levring, Y. Tanaka (Ed.), *Marine algae in pharmaceutical science* (293–302. ss.). Berlin: De Gruyter. <https://doi.org/10.1515/9783110882049>
- Fimbres-Olivarría, D., López-Elías, J. A., Carvajal-Millán, E., Márquez-Escalante, J. A., Martínez-Córdova, L. R., Miranda-Baeza, A., ... & Brown-Bojórquez, F. (2016). *Navicula* sp. sulfated polysaccharide gels induced by Fe (III): rheology and microstructure. *International journal of molecular sciences*, 17(8), 1238.
- Elion, G. B., Furman, P. A., Fyfe, J. A., de Miranda, P., Beauchamp, L., & Schaeffer, H. J. (1977). Selectivity of action of an antiherpetic agent, 9-(2-hydroxyethoxymethyl) guanine. *Proceedings of the National Academy of Sciences of the United States of America*, 74(12), 5716–5720.
- Fox, F. (2022). *Spirulina meneghiniana* cf [photographer]. Micro Foto. Retrieval date: 02/12/2022. <https://www.mikro-foto.de/algen-2.html>
- Fox, F. (2022). *Haematococcus* sp cf [photographer]. Micro Foto. Retrieval date: 02/12/2022. <https://mikro-foto.de/algen-10.html>
- Gellatly, S. L., & Hancock, R. E. (2013). *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathogens and disease*, 67(3), 159–173.

- Gerasimenko, N. I., Busarova, N. G., & Moiseenko, O. P. (2010). Age-dependent changes in the content of lipids, fatty acids, and pigments in brown alga *Costaria costata*. *Russian Journal of Plant Physiology*, 57(1), 62–68.
- Gerchman, Y., Vasker, B., Tavasi, M., Mishaël, Y., Kinel-Tahan, Y., & Yehoshua, Y. (2017). Effective harvesting of microalgae: Comparison of different polymeric flocculants. *Bioresource Technology*, 228, 141–146.
- Gerde, J. A., Wang, T., Yao, L., Jung, S., Johnson, L. A., & Lamsal, B. (2013). Optimizing protein isolation from defatted and non-defatted *Nannochloropsis* microalgae biomass. *Algal Research*, 2(2), 145–153.
- Ghasemi, Y., Tabatabaei Yazdi, M., Shafiee, A., Amini, M., Shokravi, S., & Zarrini, G. (2004). Parsiguine, a novel antimicrobial substance from *Fischerella ambigua*. *Pharmaceutical Biology*, 42(4–5), 318–322.
- Green, C. E. L., Freeman, D., Kuipers, E., Bebbington, P., Fowler, D., Dunn, G., & Garety, P. A. (2008). Measuring ideas of persecution and social reference: The Green et al. Paranoid Thought Scales (GPTS). *Psychological Medicine*, 38(1), 101–111.
- Guschina, I. A., & Harwood, J. L. (2006). Lipids and lipid metabolism in eukaryotic algae. *Progress in Lipid Research*, 45(2), 160–186.
- Hac Isa, M., Metin, C., Ercan, E., & Alparslan, Y. (2022). Effect of different cell disruption methods on lipid yield of *Schizochytrium* sp. *Journal of the American Oil Chemists' Society*, 99(2), 129–139.
- Hakim, A. R. (2012). The potential of heterotrophic microalgae (*Schizochytrium* sp.) as a source of DHA. *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 7(1), 29–38.
- Hall-Stoodley, L., & Stoodley, P. (2009). Evolving concepts in biofilm infections. *Cellular Microbiology*, 11(7), 1034–1043.
- Handayani, N. A., Ariyanti, D., & Hadiyanto, H. (2011). Potential production of polyunsaturated fatty acids from microalgae. *International Journal of Science and Engineering*, 2(1), 13–16.

- Harun, R., Singh, M., Forde, G. M., & Danquah, M. K. (2010). Bioprocess engineering of microalgae to produce a variety of consumer products. *Renewable and Sustainable Energy Reviews*, 14(3), 1037–1047.
- Herrero, M., Ibáñez, E., Cifuentes, A., Reglero, G., & Santoyo, S. (2006). *Dunaliella salina* microalga pressurized liquid extracts as potential antimicrobials. *Journal of food protection*, 69(10), 2471-2477.
- Hossain, Z., Kurihara, H., Hosokawa, M., & Takahashf, K. (2005). Growth inhibition and induction of differentiation and apoptosis mediated by sodium butyrate in Caco-2 cells with algal glycolipids. *In vitro cellular & developmental biology. Animal*, 154-159.
- Jako, C., Kumar, A., Wei, Y., Zou, J., Barton, D. L., Giblin, E. M., ... & Taylor, D. C. (2001). Seed-specific over-expression of an Arabidopsis cDNA encoding a diacylglycerol acyltransferase enhances seed oil content and seed weight. *Plant physiology*, 126(2), 861-874.
- Jakobsen, A. N., Aasen, I. M., Josefsen, K. D., & Strøm, A. R. (2008). Accumulation of docosaheptaenoic acid-rich lipid in thraustochytrid *Aurantiochytrium* sp. strain T66: Effects of N and P starvation and O₂ limitation. *Applied Microbiology and Biotechnology*, 80(2), 297–306.
- Ji, B., Zhang, W., Zhang, N., Wang, J., Lutz, G. A., & Liu, T. (2014). Biofilm cultivation of the oleaginous microalgae *Pseudochlorococcum* sp. *Bioprocess and Biosystems Engineering*, 37(7), 1369–1375.
- Kang, C. D., Han, S. J., Choi, S. P., & Sim, S. J. (2010). Fed-batch culture of astaxanthin-rich *Haematococcus pluvialis* by exponential nutrient feeding and stepwise light supplementation. *Bioprocess and Biosystems Engineering*, 33(1), 133–139.
- Katircioğlu, H., Beyatli, Y., Aslim, B., Yüksesdag, Z., & Atici, T. (2006). Screening for antimicrobial agent production of some freshwater. *The internet journal of microbiology*, 2(2).
- Kaur, G., Sarkar, N., Bhatnagar, S., Kumar, S., Raptap, C. C., Bhan, M. K., & Mehra, N. K. (2002). Pediatric celiac disease in India is associated with multiple DR3-DQ2 haplotypes. *Human Immunology*, 63(8), 677–682.

- Kellam, S. J., & Walker, J. M. (1989). Antibacterial activity from marine microalgae in laboratory culture. *British Phycolological Journal*, 24(2), 191–194.
- Kreth, J., Zhang, Y., & Herzberg, M. C. (2008). Streptococcal antagonism in oral biofilms: *Streptococcus sanguinis* and *Streptococcus gordonii* interference with *Streptococcus mutans*. *Journal of Bacteriology*, 190(13), 4632–4640.
- Lee, K. W., & Lip, G. Y. H. (2003). The role of omega-3 fatty acids in the secondary prevention of cardiovascular disease. *Qjm*, 96(7), 465–480.
- Lim, G. P., Calon, F., Morihara, T., Yang, F., Teter, B., Ubeda, O., Salem, N., Frautschy, S. A., & Cole, G. M. (2005). A diet enriched with the omega-3 fatty acid docosahexaenoic acid reduces amyloid burden in an aged Alzheimer mouse model. *Journal of Neuroscience*, 25(12), 3032–3040.
- Lima-Filho, J. V. M., Carvalho, A. F. F. U., Freitas, S. M., & Melo, V. M. M. (2002). Antibacterial activity of extracts of six macroalgae from the Northeastern Brazilian coast. *Brazilian Journal of Microbiology*, 33(4), 311–313.
- Lyczak, J. B., Cannon, C. L., & Pier, G. B. (2000). Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes and infection*, 2(9), 1051–1060.
- Mariscal, A., Carnero-Varo, M., Gutierrez-Bedmar, M., Garcia-Rodriguez, A., & Fernandez-Crehuet, J. (2007). A fluorescent method for assessing the antimicrobial efficacy of disinfectant against *Escherichia coli* ATCC 35218 biofilm. *Applied microbiology and biotechnology*, 77(1), 233–240.
- Markou, G., Depraetere, O., Vandamme, D., & Muylaert, K. (2015). Cultivation of *Chlorella vulgaris* and *Arthrospira platensis* with recovered phosphorus from wastewater by means of zeolite sorption. *International Journal of Molecular Sciences*, 16(2), 4250–4264.
- Mattson, F. H., & Grundy, S. M. (1985). Comparison of effects of dietary saturated, monounsaturated, and polyunsaturated fatty acids on plasma lipids and lipoproteins in man. *Journal of Lipid Research*, 26(2), 194–202.
- Mayer, A. M., & Hamann, M. T. (2005). Marine pharmacology in 2001–2002: Marine compounds with anthelmintic, antibacterial, anticoagulant, antidiabetic, antifungal, anti-inflammatory, antimalarial, antiplatelet, antiprotozoal, antituberculosis, and antiviral activities; affecting the cardiovascular,

immune and nervous systems and other miscellaneous mechanisms of action. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 140(3-4), 265-286.

Maltsev, Y. (2022). Unicellular green algae with large cells[photographer]. Shutter stock. Retrieval date: 02/12/2022. <https://www.shutterstock.com/image-photo/unicellular-green-algae-large-cells-part-1543549226>

McCann, M. C., Buckeridge, M. S., & Carpita, N. C. (2014). *Plants and BioEnergy*. New York, NY: Springer.

Mendes, R. L., Nobre, B. P., Cardoso, M. T., Pereira, A. P., & Palavra, A. F. (2003). Supercritical carbon dioxide extraction of compounds with pharmaceutical importance from microalgae. *Inorganica Chimica Acta*, 356, 328–334.

Merritt, J. H., Kadouri, D. E., & O'Toole, G. A. (2011). Growing and analyzing static biofilms. *Current protocols in microbiology*, 22(1), 1B-1.

Metwalli, K. H., Khan, S. A., Krom, B. P., & Jabra-Rizk, M. A. (2013). *Streptococcus mutans*, *Candida albicans*, and the human mouth: a sticky situation. *PLoS pathogens*, 9(10), e1003616.

Mobin, S. M., Kanai, K., & Yoshikoshi, K. (2001). Effects of feeding levels on the growth and survival of larval and juvenile Japanese flounder *Paralichthys olivaceus*. *Aquaculture Science*, 49(2), 207-218.

Mooij, P. R., de Jongh, L. D., van Loosdrecht, M. C. M., & Kleerebezem, R. (2016). Influence of silicate on enrichment of highly productive microalgae from a mixed culture. *Journal of Applied Phycology*, 28(3), 1453–1457.

Mozaffarian, D., & Wu, J. H. Y. (2011). Omega-3 fatty acids and cardiovascular disease: Effects on risk factors, molecular pathways, and clinical events. *Journal of the American College of Cardiology*, 58(20), 2047–2067.

Mühlroth, A., Li, K., Røkke, G., Winge, P., Olsen, Y., Hohmann-Marriott, M. F., Vadstein, O., & Bones, A. M. (2013). Pathways of lipid metabolism in marine algae, co-expression network, bottlenecks and candidate genes for enhanced production of EPA and DHA in species of chromista. *Marine Drugs*, 11(11), 4662–4697.

- Mullen, A., Loscher, C. E., & Roche, H. M. (2010). Anti-inflammatory effects of EPA and DHA are dependent upon time and dose-response elements associated with LPS stimulation in THP-1-derived macrophages. *Journal of Nutritional Biochemistry*, 21(5), 444–450.
- Nag, M., Lahiri, D., Dey, A., Sarkar, T., Joshi, S., & Ray, R. R. (2022). Evaluation of algal active compounds as potent antibiofilm agent. *Journal of Basic Microbiology*, 62(9), 1098-1109.
- Nichols, P. D., Petrie, J., & Singh, S. (2010). Long-chain omega-3 oils-an update on sustainable sources. *Nutrients*, 2(6), 572–585.
- Nichols, P. D., McManus, A., Krail, K., Sinclair, A. J., & Miller, M. (2014). Recent advances in omega-3: Health benefits, Sources, Products and bioavailability. *Nutrients*, 6(9), 3727–3733.
- Ohta, S., Shiomi, Y., Kawashima, A., Aozasa, O., Nakao, T., Nagate, T., ... & Miyata, H. (1995). Antibiotic effect of linolenic acid from *Chlorococcum* strain HS-101 and *Dunaliella primolecta* on methicillin-resistant *Staphylococcus aureus*. *Journal of applied phycology*, 7(2), 121-127.
- Ohta, S., Ono, F., Shiomi, Y., Nakao, T., Aozasa, O., Nagate, T., Kitamura, K., Yamaguchi, S., Nishi, M., & Miyata, H. (1998). Anti-herpes simplex virus substances produced by the marine green alga, *Dunaliella primolecta*. *Journal of Applied Phycology*, 10(4), 349–356.
- P. Desbois, A. (2012). Potential Applications of Antimicrobial Fatty Acids in Medicine, Agriculture and Other Industries. *Recent Patents on Anti-Infective Drug Discovery*, 7(2), 111–122.
- Pal, D., Khozin-Goldberg, I., Cohen, Z., & Boussiba, S. (2011). The effect of light, salinity, and nitrogen availability on lipid production by *Nannochloropsis* sp. *Applied Microbiology and Biotechnology*, 90(4), 1429–1441.
- Paniagua-Michel, J. D. J., Morales-Guerrero, E., & Olmos Soto, J. (2015). Microalgal biotechnology: biofuels and bioproducts. K. Se-Kwon (Ed.), *Springer handbook of marine biotechnology* (1355-1370. ss.). Berlin, Heidelberg: Springer. https://doi.org/10.1007/978-3-642-53971-8_62
- Parka, Y., & Harris, W. (2002). EPA, but not DHA, decreases mean platelet volume in normal subjects. *Lipids*, 37(10), 941–946.

- Parsaeimehr, A., Sun, Z., Dou, X., & Chen, Y. F. (2015). Simultaneous improvement in production of microalgal biodiesel and high-value alpha-linolenic acid by a single regulator acetylcholine. *Biotechnology for Biofuels*, 8(1), 1–10.
- Parsaeimehr, A., & Lutz, G. A. (2016). Algae as a novel source of antimicrobial compounds: current and future perspectives. K. Kon, M. Rai (Ed.), *Antibiotic Resistance* (377-413. ss.). <https://doi.org/10.1016/B978-0-12-803642-6.00018-6>
- Patel, D. S., Pendrill, R., Mallajosyula, S. S., Widmalm, G., & MacKerell, A. D. (2014). Conformational properties of α - Or β -(1→6)-linked oligosaccharides: Hamiltonian replica exchange MD simulations and NMR experiments. *Journal of Physical Chemistry B*, 118(11), 2851–2871.
- Petrie, J. R., Shrestha, P., Mansour, M. P., Nichols, P. D., Liu, Q., & Singh, S. P. (2010). Metabolic engineering of omega-3 long-chain polyunsaturated fatty acids in plants using an acyl-CoA Δ 6-desaturase with ω 3-preference from the marine microalga *Micromonas pusilla*. *Metabolic Engineering*, 12(3), 233–240.
- Peltomaa, E., Johnson, M. D., & Taipale, S. J. (2018). Marine cryptophytes are great sources of EPA and DHA. *Marine Drugs*, 16(1), 3.
- Perez-Garcia, O., Escalante, F. M. E., de-Bashan, L. E., & Bashan, Y. (2011). Heterotrophic cultures of microalgae: Metabolism and potential products. *Water Research*, 45(1), 11–36.
- Pericolini, E., Colombari, B., Ferretti, G., Iseppi, R., Ardizzoni, A., Girardis, M., ... & Blasi, E. (2018). Real-time monitoring of *Pseudomonas aeruginosa* biofilm formation on endotracheal tubes in vitro. *BMC microbiology*, 18(1), 1-10.
- Phiri, R. M. (2021). Determination of cyanobacteria diversity from hot springs around Aydin province by culture independent methods. (Master dissertation). Retrieved from <https://tez.yok.gov.tr/UlusalTezMerkezi/giris.jsp> (FEF- 19020).
- Plaza, M., Santoyo, S., Jaime, L., García-Blairsy Reina, G., Herrero, M., Señoráns, F. J., & Ibáñez, E. (2010). Screening for bioactive compounds from algae. *Journal of Pharmaceutical and Biomedical Analysis*, 51(2), 450–455.

- Rath, P., Behura, N., Sahoo, S., Panda, P., Mandal, K., & Panigrahi, P. (2015). Amelioration of heat stress for poultry welfare: a strategic approach. *International Journal of Livestock Research*, 5(3), 1-9.
- Ratledge, C. (2004). Fatty acid biosynthesis in microorganisms being used for Single Cell Oil production. *Biochimie*, 86(11), 807–815.
- Rappe, M. (2019). *Dunaliella* saltwater algae[photographer]. Cals news. Retrieval date: 24.12.2022. <https://cals.ncsu.edu/news/salty-algae-leads-to-sweet-success/>.
- Robertson, R., Guihéneuf, F., Schmid, M., Stengel, D. B., Fitzgerald, G., Ross, P., & Stanton, C. (2013). Algae-derived polyunsaturated fatty acids: implications for human health. A. Catala (Ed.), *Polyunsaturated fatty acids: sources, antioxidant properties and health benefits* (45-99. ss.). New York: Nova Sciences Publishers, Inc.
- Rodríguez-Meizoso, I., Jaime, L., Santoyo, S., Señoráns, F. J., Cifuentes, A., & Ibáñez, E. (2010). Subcritical water extraction and characterization of bioactive compounds from *Haematococcus pluvialis* microalga. *Journal of pharmaceutical and biomedical analysis*, 51(2), 456-463.
- Romeo, T. (2008). *Bacterial Biofilms- Current topics in microbiology and immunology*. New York: Springer Publishers.
- Ryckebosch, E., Bruneel, C., Termote-Verhalle, R., Goiris, K., Muylaert, K., & Foubert, I. (2014). Nutritional evaluation of microalgae oils rich in omega-3 long chain polyunsaturated fatty acids as an alternative for fish oil. *Food Chemistry*, 160, 393-400.
- Ryder, C., Byrd, M., & Wozniak, D. J. (2007). Role of polysaccharides in *Pseudomonas aeruginosa* biofilm development. *Current opinion in microbiology*, 10(6), 644-648.
- Santoyo, S., Rodríguez-Meizoso, I., Cifuentes, A., Jaime, L., Reina, G. G. B., Señorans, F. J., & Ibáñez, E. (2009). Green processes based on the extraction with pressurized fluids to obtain potent antimicrobials from *Haematococcus pluvialis* microalgae. *LWT-Food Science and Technology*, 42(7), 1213-1218.

- Santoyo, S., Jaime, L., Plaza, M., Herrero, M., Rodriguez-Meizoso, I., Ibañez, E., & Reglero, G. (2012). Antiviral compounds obtained from microalgae commonly used as carotenoid sources. *Journal of applied phycology*, 24(4), 731-741.
- Sampathkumar, S. J., Srivastava, P., Ramachandran, S., Sivashanmugam, K., & Gothandam, K. M. (2019). Lutein: A potential antibiofilm and antiquorum sensing molecule from green microalga *Chlorella pyrenoidosa*. *Microbial pathogenesis*, 135, 103658.
- Saravanakumar, D. E. M., Folb, P. I., Campbell, B. W., & Smith, P. (2008). Antimycobacterial Activity of the Red Alga *Polysiphonia virgata*. *Pharmaceutical Biology*, 46(4), 254-260.
- Schulze, C., Wetzel, M., Reinhardt, J., Schmidt, M., Felten, L., & Mundt, S. (2016). Screening of microalgae for primary metabolites including β -glucans and the influence of nitrate starvation and irradiance on β -glucan production. *Journal of Applied Phycology*, 28(5), 2719–2725.
- Shah, M. M. R., Liang, Y., Cheng, J. J., & Daroch, M. (2016). Astaxanthin-producing green microalga *Haematococcus pluvialis*: from single cell to high value commercial products. *Frontiers in plant science*, 7, 531.
- Simopoulos, A. P. (2002). Omega-3 fatty acids in inflammation and autoimmune diseases. *Journal of the American College of Nutrition*, 21(6), 495–505.
- Smith, V. J., Desbois, A. P., & Dyrinda, E. A. (2010). Conventional and unconventional antimicrobials from fish, marine invertebrates and micro-algae. *Marine Drugs*, 8(4), 1213–1262.
- Stabili, L., Acquaviva, M. I., Biandolino, F., Cavallo, R. A., De Pascali, S. A., Fanizzi, F. P., ... & Cecere, E. (2012). The lipidic extract of the seaweed *Gracilariopsis longissima* (Rhodophyta, Gracilariales): a potential resource for biotechnological purposes?. *New Biotechnology*, 29(3), 443-450.
- Strunecký, O., Kopejtká, K., Goecke, F., Tomasch, J., Lukavský, J., Neori, A., Kahl, S., Pieper, D. H., Pilarski, P., Kaftan, D., & Koblížek, M. (2019). High diversity of thermophilic cyanobacteria in Rupite hot spring identified by microscopy, cultivation, single-cell PCR and amplicon sequencing. *Extremophiles*, 23(1), 35–48.

- Sun, J., & Song, X. (2011). Assessment of antimicrobial susceptibility of *Enterococcus faecalis* isolated from chronic periodontitis in biofilm versus planktonic phase. *Journal of periodontology*, 82(4), 626-631.
- Sun, M., Zhou, Z., Dong, J., Zhang, J., Xia, Y., & Shu, R. (2016). Antibacterial and antibiofilm activities of docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) against periodontopathic bacteria. *Microbial Pathogenesis*, 99, 196–203.
- Surette, M. E. (2008). The science behind dietary omega-3 fatty acids. *Cmaj*, 178(2), 177-180.
- Swanson, D., Block, R., & Mousa, S. A. (2012). Omega-3 fatty acids EPA and DHA: Health benefits throughout life. *Advances in Nutrition*, 3(1), 1–7.
- Thormar, H. (2010). *Lipids and Essential Oils as Antimicrobial Agents*. Chichester, UK : John Wiley & Sons Ltd. Publication.
- Thongdumhyu, P. (2018). *Chlorella* under the microscope in Lab [photographer]. Dreams time. Retrieval date: 18.12.2022. <https://www.dreamstime.com/Chlorella-under-microscope-lab-education-image158565381>.
- Tunçer, S., & Karaçam, S. (2020). Cell-free supernatant of *Streptococcus salivarius* M18 impairs the pathogenic properties of *Pseudomonas aeruginosa* and *Klebsiella pneumonia*. *Archives of Microbiology*, 202(10), 2825-2840.
- Ursu, A. V., Marcati, A., Sayd, T., Sante-Lhoutellier, V., Djelveh, G., & Michaud, P. (2014). Extraction, fractionation and functional properties of proteins from the microalgae *Chlorella vulgaris*. *Bioresource Technology*, 157, 134–139.
- Uzundağ, D. (2020). Biofilm Formation Mechanism in Fungi. *Anatolian Journal of Botany*, 4(2), 116–120.
- Vairappan, C. S. (2003). Potent antibacterial activity of halogenated metabolites from Malaysian red algae, *Laurencia majuscula* (Rhodomelaceae, Ceramiales). *Biomolecular engineering*, 20(4-6), 255-259.

- Vardar, N. (2016). Optimization of cultivation conditions and engineering by random mutagenesis for high lipid production in *Schizochytrium* sp. S31. (Doctoral dissertation). Retrieved from National Thesis Center (450950).
- Vardar, N., Ayhan, Y., Yelboğa, E., Bahar, B., Tüter, M., & Karagüler, N. G. (2016). Optimization of cultivation and supercritical fluid lipid extraction conditions of *Schizochytrium* sp. S31. *Int J Eng Sci*, 2, 13-24.
- Vardar-Yel, N., Yelboğa, E., Tüter, M., & Karagüler, N. G. (2017). Comparison of cell disruption and lipid extraction methods for improving lipid content of *Schizochytrium* sp. S31. *J Mol Biol Biotechnol*, 1(1), 9-12.
- Van Houdt, R., & Michiels, C. W. (2005). Role of bacterial cell surface structures in *Escherichia coli* biofilm formation. *Research in Microbiology*, 156(5–6), 626–633.
- Veetil, S. R., Sunil, E. A., Mukunda, A., Mohan, A., John, S., & Pynadath, M. K. (2022). Anticancer effect of Piper betle leaf extract on KB cell lines-an in vitro study. *Oral & Maxillofacial Pathology Journal*, 13(1).
- Wang, J., Sommerfeld, M. R., Lu, C., & Hu, Q. (2013). Combined effect of initial biomass density and nitrogen concentration on growth and astaxanthin production of *Haematococcus pluvialis* (Chlorophyta) in outdoor cultivation. *Algae*, 28(2), 193–202.
- Wang, Y., Tibbetts, S. M., & McGinn, P. J. (2021). Microalgae as sources of high-quality protein for human food and protein supplements. *Foods*, 10(12), 3002.
- Wannigama, D. L., Hurst, C., Hongsing, P., Pearson, L., Saethang, T., Chantaravisoot, N., ... & Chatsuwan, T. (2020). A rapid and simple method for routine determination of antibiotic sensitivity to biofilm populations of *Pseudomonas aeruginosa*. *Annals of clinical microbiology and antimicrobials*, 19(1), 1-8.
- Ward, O. P., & Singh, A. (2005). Omega-3/6 fatty acids: alternative sources of production. *Process biochemistry*, 40(12), 3627-3652.

- Wey, S. B., Mori, M., Pfaller, M. A., Woolson, R. F., & Wenzel, R. P. (1988). Hospital-acquired candidemia: the attributable mortality and excess length of stay. *Archives of internal medicine*, 148(12), 2642-2645.
- Whitton, B. A. (2012). *Ecology of cyanobacteria II: Their diversity in space and time*. New York: Springer Publishers.
- Wille, J. J., & Kydonieus, A. J. S. P. (2003). Palmitoleic acid isomer (C16: 1Δ6) in human skin sebum is effective against gram-positive bacteria. *Skin Pharmacology and Physiology*, 16(3), 176-187.
- Xin, Y., Shen, C., She, Y., Chen, H., Wang, C., Wei, L., Yoon, K., Han, D., Hu, Q., & Xu, J. (2019). Biosynthesis of Triacylglycerol Molecules with a Tailored PUFA Profile in Industrial Microalgae. *Molecular Plant*, 12(4), 474–488.
- Yi, Z., Yin-shan, C., & Hai-sheng, L. (2001). Screening for antibacterial and antifungal activities in some marine algae from the Fujian coast of China with three different solvents. *Chinese Journal of Oceanology and Limnology*, 19(4), 327–331.
- Zhang, H., Zhang, L., Peng, L. J., Dong, X. W., Wu, D., Wu, V. C. H., & Feng, F. Q. (2012). Quantitative structure-activity relationships of antimicrobial fatty acids and derivatives against *Staphylococcus aureus*. *Journal of Zhejiang University: Science B*, 13(2), 83–93.
- Zhang, S., Yang, G., Ye, Q., Wu, Q., Zhang, J., & Huang, Y. (2018). Phenotypic and genotypic characterization of *Klebsiella pneumoniae* isolated from retail foods in China. *Frontiers in microbiology*, 9, 289.
- Zhou, F., Wang, D., Hu, J., Zhang, Y., Tan, B. K., & Lin, S. (2022). Control Measurements of *Escherichia coli* Biofilm: A Review. *Foods*, 11(16), 2469.
- Zhu, L. D., Hiltunen, E., Antila, E., Zhong, J. J., Yuan, Z. H., & Wang, Z. M. (2014). Microalgal biofuels: Flexible bioenergies for sustainable development. *Renewable and Sustainable Energy Reviews*, 30, 1035-1046.

APPENDIX A

A.1 Chemical Characteristics of the of *Schizochytrium* sp. S31 Dry Biomass

A.2 Total Amino Acid and Vitamin Content of the Dry Biomass

A.3 Gas Chromatography Analysis Report



A.1 Chemical Characteristics of the of *Schizochytrium* sp. S31 Dry Biomass



Rapor No (Report Number) : 2112367
Revizyon No (Revision No) : 00 Rapor Yayın Tarihi (Date of Issue) : 29.09.2021

Analiz Amacı (Purpose of Analysis)		Özel İstek (Private Request)					
Numuneyi Gönderen / Adresi (Sample Sender / Address)		MARİN BİYOTEKNOLOJİ ÜRÜNLERİ VE GIDA SAN. TİC. LTD. ŞTİ. / ATA OSA MAH.					
Laboratuvara Kabul Tarihi (Laboratory Acceptance Date)		22.09.2021 / 17:12					
Analiz Başlangıç ve Bitiş Tarihi (Start and End of Analysis)		22.09.2021 / 20.09.2021					
Numune (Sample)		Gıda (FOOD)					
Cinsi (Type)		Gıda (FOOD)					
Adı (Name)		MİKROALG					
Markası (Brand)		-					
Ambalajı (Packaging)		Kilitli Poşet (Locked Bag)					
Miktar / Adedi (Amount/Pieces)		200 g					
Parti No/ Serisi / Mühür No (Lot/Series/Seal No)		-					
Üretim ve Son Kullanma Tarihi (Production and Expire Date)		-					
Açıklama/ Revizyon Nedeni (Explanation / Reason for Revision)		-					
Analizler (Analyses)	Sonuç (Result)	OB (MU) (OB) (RN)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LDK (Sol.)	DS (Evolution)	
1- * Toplam Yağ Miktarı Tayini (Determination of Total Fat And Oil) (g/100g)	5,31	0.875	-	NMKL 100			
2- * Azot (Protein) Miktarı Tayini (Determination of Nitrogen (Protein) Quantity) (g/100g(Nx6,25))	44,62	0.056	-	TS ISO 1871, TS 1748 ISO 937			

LOQ: Ölçüm limiti (Limit of Quantification) ; OB: Belirli kasanın (Recovery); OB: Ölçüm belirsizliği (MU: Measurement Uncertainty); LDK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirmeye Sonuçta (Evaluation) ; U: Uygun (Confirmed); UO: Uygun Değil (Not Confirmed); D: Değerlendirmeye Yapılmadı (No Evaluation)

Birim Sorumlusu/ Department Responsible

Nurcan KOC
Fiziksel Analiz Birim Sorumlusu
Head of Physical Analysis Dept.

Serkan TEKİN
Kimyasal Analiz Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKBUĞAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and Reporting Dept.

Tasdik Olunur (Approved by)
29.09.2021

Dr. Veyisel Umut ÇİLDIK
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/03/20.06.2021

Sayfa (Page) 1 / 4

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Mersis No : 0125 0259 0550 0015

Karacaöğlen Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 493 00 81
T | +90 232 493 00 57
F | +90 232 472 13 22



Tarih
18.09.2021
AB-0705-7

2112367

09-21

T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORY)
MUAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Rapor No (Report Number) : 2112367
Revizyon No (Revision No) : 00

Rapor Yayın Tarihi (Date of Issue): 29.09.2021

Analizler (Analyses)	Sonuç (Result)	Ölçü (MU)	GR(%) (%)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LİMİ (Lim)	DE (Evaluation)
3- * Enerji tayini (kcal/100g) (Determination of Energy (kcal / 100g)) (kcal/100g)	374,95	35,22	-	-	26.01.2017 25960 Sayılı Resmî Gazete Türk Gıda Kodeksi Gıda Etiketleme ve Tüketici Bilgilendirme Yönetmeliği		
4- * Karbonhidrat Tayini (Determination of Carbohydrate) (g/100g)	37,17	3,591	-	-	26.01.2017 25960 Sayılı Resmî Gazete Türk Gıda Kodeksi Gıda Etiketleme ve Tüketici Bilgilendirme Yönetmeliği		
5- * Enerji tayini (kJ/100g) (Determination of Energy (kJ/100g)) (kJ/100g)	1568,79	151,545	-	-	26.01.2017 25960 Sayılı Resmî Gazete Türk Gıda Kodeksi Gıda Etiketleme ve Tüketici Bilgilendirme Yönetmeliği		

LOQ: Ölçüm limiti (Limit of Quantitation), GR: Belirli Kaan oran (Ratio), DE: Değerlendirme (Assessment), LİMİ: Limit Değer Kaynağı (Source of Limit)
DE: Değerlendirme sonucu (Evaluation), U: Uygun (Conformed), GÜ: Uygun Değil (Not Conformed), DE: Değerlendirme Yapılmadı (Not Evaluation)

Birim Sorumlusu/ Department Responsible

Nurdan KOC
Fiziksel An.Lab.Birim Sorumlusu
Head of Physical Analysis Dept.

Serkan TEKİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKGÜDAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tasdik Olunur (Approved by)
29.09.2021

Dr. Veyisel Umut ÇELİK
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/03/03.06.2021

Sayfa (Page) 2 / 4

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Mersis No : 0125.0256.0550.0005

Karacaoğlu Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 492 00 81
T | +90 232 492 00 57
F | +90 232 472 13 22



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MÜHÜR VE ANALİZ RAPORU
(ANALYSIS REPORT)

AB-0705-T
21.12.2021
09-21

Rapor No (Report Number) : 2112367
Revizyon No (Revision No) : 00

Rapor Yayın Tarihi (Date of Issue): 29.09.2021

Analizler (Analyses)	Sonuç (Result)	OB (MO)	GR[N] (RN)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LDE (Sol)	DE (Evaluation)
6- * Rutubet ve/veya Uçucu Madde, Kurutma Kaybı, Nem Miktari Tayini (Determination of Moisture and / or Volatile Substance, Drying loss, Moisture content) (g/100g)	7,19	0.829	-	-	TS EN ISO 712		
7- * Kül / Ham Kül Tayini (Total ash) (g/100g)	5,71	0.818	-	-	TS EN ISO 2171		

LOQ: Ölçme sınırı (Limit of Quantitation) ; OB: Geni Kapanım (Recovery); GR: Ölçüm Belirsizliği (Std. Measurement uncertainty); LDE: Limit Değer Kaynağı (Source of Limit)
DE: Değerlendirme Sonucu (Evaluation) ; U: Uygun (Conformed); SU: Uygun Değil (Not Conformed); DE: Değerlendirme Yapılmadı (Not Evaluated)

Birim Sorumlusu/ Department Responsible

Nurdan KOC
Fiziksel An.Lab.Birim Sorumlusu
Head of Physical Analysis Dept.

Serkan TEKİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKBUĞAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tamam Olunur (Approved by)
29.09.2021

Dr. Veyuel Umut ÇİLDİR
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/01/30.06.2021

Sayfa (Page) 3 / 4

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Mersis No : 0125 0256 0550 00015

Karacaoğlu Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 492 00 81
T | +90 232 492 00 57
F | +90 232 472 13 22

A.2 Total Amino Acid and Vitamin Content of the Dry Biomass



Amino Acid	g/100g protein
Aspartik asit	10.5 ± 0.09
Treonin	5.24 ± 0.12
Serin	5.08 ± 0.07
Glutamik asit	10.7 ± 0.03
Prolin	6.45 ± 0.26
Glisin	5.13 ± 0.07
Alanin	8.44 ± 0.19
Valin	6.44 ± 0.12
Methionin	1.52 ± 0.13
İzolösin	5.01 ± 0.14
Lösin	6.84 ± 0.17
Tirosin	5.21 ± 0.21
Fenilalanin	4.26 ± 0.04
Histidin	5.02 ± 0.10
Lisin	5.63 ± 0.08
Arjinin	8.25 ± 0.26

info@marinbio.com.tr • www.marinbio.com.tr • Tel: 256 231 10 25 • Fax: 256 231 10 26
Ata Mh. 769 Sk. 97/1 Efeler, AYDIN




Marin Biyoteknoloji Ürünleri ve Gıda San. Tic. Ltd. Şti.

Vitamin	Concentration
Demir, mg/kg	869.51 ± 38.96
Niyasin, mg/kg	245 ± 16.58
Potasyum, mg/kg	1360 ± 241.74
Kalsiyum, mg/kg	100.08 ± 9.62
Lutein, mg/kg	368.65 ± 56.72
A vitamini, IU/kg	216.41 ± 72.13
B ₁ vitamini, mg/kg	4.48 ± 0.12
B ₂ vitamini, mg/kg	14.26 ± 3.04
B ₆ vitamini, mg/kg	9.62 ± 2.84
C vitamini, mg/kg	74.16 ± 1.85
B ₁₂ vitamini, µg/kg	725.17 ± 156.20

info@marinbio.com.tr • www.marinbio.com.tr • Tel: 256 231 10 25 • Fax: 256 231 10 26
Ata Mh. 769 Sk. 97/1 Efeler, AYDIN

A.3 Gas Chromatography Analysis Report



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MÜAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Yayın Tarihi (Date of Issue): 20.10.2021

Analiz Amacı (Purpose of Analysis)	: Özel İstek (Private Request)						
Numaralı Gönderen (Sample Sender)	: Atınbağ Üniversitesi						
Laboratuvara Kabul Tarihi (Laboratory Acceptance Date)	: 18.10.2021 / 13:26						
Analiz Başlangıç ve Bitiş Tarihi (Start and End of Analysis)	: 18.10.2021 / 20.10.2021						
Örnekler (Sample/s)	: Bitkisel / Hayvansal Yağ						
Örnek Türü (Type)	: Bitkisel / Hayvansal Yağ						
Adı (Name)	: MİRRDALG YAĞI						
Markası (Brand)	: -						
Ambalajı (Package)	: Plastik Kap (Plastic package)						
Miktar / Adedi (Amount/Pieces)	: 5 ml.						
Parti No/ Seri / Mühür No (Lot/Seria/Seal No)	: 10/2021						
Üretim ve Son Kullanma Tarihi (Production and Expiry Date)	: -						
Açıklama/ Revizyon Nedeni (Explanation / Reason for Revision)	: -						
Analizler (Analyses)	Sonuç (Result)	Ölç (MU) (MU)	GR(%) (GR)	LDK (LDK)	Analiz Metodu (Analysis Method)	LDK (LDK)	DS (Evaluation)
1. - Yağ Asitleri Kompozisyonu Tayini (Fatty Acids Composition) (%)	-	-	-	-	COI/T.20/Doc No:33	-	-
- Butirik Asit (C4:0) (-Butyric acid C4:0) (%)	Tespit Edilmedi (Not Detected)	0	-	-	-	-	-
- Alfa Linolenik Asit (C18:3 (n3)) (-Alfa Linolenic Acid (C18:3 (n3))) (%)	0,84	0,044	-	-	-	-	-

LDK: Ölçüm Limiti (Limit of Quantification), G.C. Geri Kazanım (Recovery), ÖB: Ölçüm Belirsizliği (MU) Measurement Uncertainty, LDK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation) : U: Uygun (Confirmed), LD: Uygun Değil (Not Confirmed), Dİ: Değerlendirme Yapılamadı (No Evaluation)

Birim Sorumlusu/ Department Responsible

Serkan TEKİN
Kimyasal An. Lab. Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKRUÖNE
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tesdik Olunur (Approved by)
20.10.2021

Dr. Veyisel Unut ÇELEME
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/03/10.06.2021

Sayfa (Page) 1 / 6

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Menis No : 05.25.0259.0550.0015

Karacaöğlan Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 462 00 81
T | +90 232 462 00 57
F | +90 232 472 13 22



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MUAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Test
T.C. EN GIDA KONTROL
AB-0785-T
AB-0785-T
2113737
10-21

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Yayımlı Tarihi (Date of Issue): 20.10.2021

Analizler (Analyses)	Sonuç (Result)	ÖB (ML)	GR(%) (%)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LOK (Sol.)	DS (Evaluation)
* -Araçidik Asit (C20:0) (- Arachidic Acid (C20:0)) (%)	0,41	0,050	-	-			
* -Araçidonic Asit (C20:4) (- Arachidonic Acid (C20:4)) (%)	1,07	0,096	-	-			
* -Behenik Asit (C22:0) (%)	0,25	0,049	-	-			
* -Cis-11 Eikosenolik Asit (C20:1) (-Cis-11 Eicosenoic Acid (C20:1)) (%)	1,15	0,129	-	-			
* -Cis-11, 14, 17 Eikosatrienolik Asit (C20:3 (n3)) (- Cis-11, 14, 17 Eicosatrienoic Acid (C20:3 (n3)) (%)	0,61	0,105	-	-			
* -Cis-8, 11, 14 Eikosatrienolik Asit (C20:3 (n6)) (-Cis-8, 11, 14 Eicosatrienoic Acid (C20:3 (n6)) (%)	Tespit Edilmedi (Not Detected)	0	-	-			
* -Cis-11, 14 Eikosadienolik Asit (C20:2) (-Cis-11, 14 Eicosadienoic Acid (C20:2)) (%)	0,17	0,023	-	-			
* -Dokosaheksanolik Asit (DHA) (C22:6) (-Docosahexanoic Acid (DHA) (C22:6)) (%)	10,47	0,550	-	-			
* -Dokosapentanolik Asit (C22:5) (-Docosapentanoic Acid (C22: 5)) (%)	1,84	0,116	-	-			

LOQ: Ölçüm Limiti (Limit of Quantification) / G.E. Geri Kazanım (Recovery) / ÖB: Ölçüm Belirsizliği (MU: Measurement Uncertainty), LOK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation) / U: Uygun (Confirmed), UY: Uygun Değil (Not Confirmed), DY: Değerlendirme Yapılamadı (No Evaluation)

Birim Sorumlusu / Department Responsible

Sehan TEKİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKSUOĞAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tasdik Olunur (Approved by)
20.10.2021

Dr. Veynel Umut ÇELİK
Laboratuvar Müdürü
Laboratory Manager


F-7.8/001/03/20.06.2021

Sayfa (Page) 2 / 6

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Menisik No : 0025.0259.0550.0015

Karacaoğlu Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 462 00 81
T | +90 232 462 00 57
F | +90 232 472 11 22



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MUAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Verim Tarihi (Date of Issue): 20.10.2021

Analizler (Analyses)	Sonuç (Result)	Ölçü (MU)	GR(%) (%)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LOK (Lot)	DS (Evaluation)
* -Eikosapentenoik Asit (EPA) (C20:5) (-Eicosapentenoic Acid (EPA) (C20:5)) (%)	10,11	0,470	-	-			
* -Gamma Linolenik Asit (C18:3 (n6)) (- Gamma Linolenic Acid (C18:3 (n6)) (%)	Tespit Edilmedi (Not Detected)	0	-	-			
* -Henekosanoik Asit (C21:0) (- Hemicosanoic Acid (C21:0)) (%)	2,93	0,135	-	-			
* -Margarik (C17:0) (- Heptadecanoic Acid (C17:0)) (%)	0,76	0,067	-	-			
* -Heptadeseenik (C17:1) (- Heptadecenoic Acid (C17:1)) (%)	0,05	0,006	-	-			
* -Kaprlik Asit (C10:0) (-Capric Acid (C10:0)) (%)	Tespit Edilmedi (Not Detected)	0	-	-			
* -Kaprlik Asit (C8:0) (-Caprylic Acid (C8:0)) (%)	Tespit Edilmedi (Not Detected)	0	-	-			
* -Kaproik Asit (C6:0) (-Caproic Acid (C6:0)) (%)	Tespit Edilmedi (Not Detected)	0	-	-			
* -Laurik Asit (C12:0) (-Lauric acid (C12:0)) (%)	0,17	0,029	-	-			

LOQ: Ölçüm Limiti (Limit of Quantification), G.E. Sani Kaşınım (Recovery), Ölç. Ölçüm Belirsizliği (MU): Measurement Uncertainty, LOK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation), U: Uygun (Confirmed), LO: Uygun Değil (Not Confirmed), EN: Değerlendirme Yapılmadı (No Evaluation)

Birim Sorumlusu/ Department Responsible

Serkan TİDİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKBUĞAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tasdik Olunur (Approved by)
20.10.2021

Dr. Veyisel Umut ÇELENKİ
Laboratuvar Müdürü
Laboratory Manager

F-7.İ/001/03/30.06.2021

Sayfa (Page) 3 / 6

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Menis No : 0625.0250.0550.0015

Karacaoğlu Mh. 6166 Sk. No:27/A
26070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 462 00 81
T | +90 232 462 00 57
F | +90 232 472 13 22



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MÜAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Test T.C. EN BS/EC 1709 AL-0190-1
AB-0765-T
2113737
10-21

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Yayın Tarihi (Date of Issue): 20.10.2021

Analizler (Analyses)	Sonuç (Result)	ÖB (M1)	GR(%) (%)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LOK (Sk.)	DS (Evaluation)
* -Lignoserik Asit (C24:0) (- Lignoseric Acid (C24:0)) (%)	0,70	0,211	-	-			
* -Erusik asit C22:1 (-Erusic Acid (C22:1)) (%)	0,11	0,009	-	-			
* -Linoleik Asit (C18:2) (-Linoleic Acid (C18:2)) (%)	1,49	0,035	-	-			
* -Miristik Asit (C14:0) (-Miristic Acid (C14:0)) (%)	7,92	1,797	-	-			
* -Miristoleik Asit (C14:1) (- Miristoleic Acid (C14:1)) (%)	0,23	0,021	-	-			
* -Nervonik Asit (C24:1) (- Nervonic Acid (C24:1)) (%)	0,33	0,059	-	-			
* -Oleik Asit (C18:1) (-Oleic Acid (C18:1)) (%)	16,80	0,339	-	-			
* -Palmirik Asit (C16:0) (- Palmitic Acid(C16:0)) (%)	14,38	1,171	-	-			
* -Pentadekanolik Asit (C15:0) (- Penta-Decanoic Acid (C15:0)) (%)	0,49	0,038	-	-			
* -Palmistoleik Asit (C16:1) (- Palmitoleic Acid (C16:1)) (%)	10,11	0,976	-	-			
* -Pentadekanolik Asit (C15:1) (- Penta-Decenoic acid (C15:1)) (%)	0,15	0,012	-	-			
* -Steirik Asit (C18:0) (-Stearic Acid (C18:0)) (%)	2,84	0,097	-	-			

LOQ: Ölçüm Limiti (Limit of Quantification) / ÖB: Ölçüm Kesimi (Reference) / GR: Ölçüm Belirsizliği (MU: Measurement Uncertainty), LOK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation) / U: Uygun (Confirmed), LOQ: Uygun Değil (Not Confirmed), DS: Değerlendirme Yapılmadı (Not Evaluation)

Birim Sorumlusu/ Department Responsible

Serkan TEĐİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKGÜDAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tasdik Olunur (Approved By)
20.10.2021

Dr. Veynel Unut ÇELEK
Laboratuvar Müdürü
Laboratory Manager

F-7.3/001/03/20.06.2021

Sayfa (Page) 4 / 6

Eurofins İzmir GıdaAnaliz
Laboratuvarları Limited Şirketi
Menis No : 0625.0259.0550.0015

Karacaoğlu Mh. 6166 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 232 460 00 81
T | +90 232 460 00 57
F | +90 232 472 13 22



Tel
+90 332 462 00 57
+90 332 462 00 57

AG-0765-T

2113737

10-21

T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MÜAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Yayımlı Tarihi (Date of Issue): 20.10.2021

Analizler (Analyses)	Sonuç (Result)	ÖB (NMI)	GR(%) (%)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LDK (Sol.)	DS (Evaluation)
* -Toplam Omega - 3 + Omega - 6 Yağ Asitleri (-Total Omega-3 + Omega-6 Fatty Acids) (%)	32,15	2,509	-	-			
* -Toplam Omega - 3 Yağ Asitleri (-Total Omega-3 Fatty Acids) (%)	27,42	2,191	-	-			
* -Toplam Omega - 6 Yağ Asitleri (-Total Omega-6 Fatty Acids) (%)	4,73	0,175	-	-			
* -Trans Linolenik Asit (IC18:3) (-Trans Linolenic Acid (IC18:3)) (%)	0,13	0,038	-	-			
* -Trans-9-Oktadekenoik Asit (Trans Oleik Asit) (IC18:1) (- Trans-9-Octadecenoic Acid (Trans Oleic Acid) (IC18:1)) (%)	0,15	0,063	-	-			
* -Trans-9,12-Oktadekadienoik Asit (Trans Linoleik Asit) (IC18:2) (-Trans-9,12- Octadecenoic Acid (Trans Linoleic Acid) (IC18:2)) (%)	0,06	0,009	-	-			
* -Trikosanoik Asit (C23:0) (- Tricosanoic Acid (C23:0)) (%)	0,75	0,065	-	-			
* -Tanımlanamayan Yağ Asitleri (Unidentified Fatty Acids) (%)	6,38	1,404	-	-			

LOQ: Ölçüm Limiti (Limit of Quantification), GR: Geri Kazanım (Recovery), ÖB: Ölçüm Belirsizliği (MI: Measurement Uncertainty), LDK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation) : U: Uygun (Confirmed), UY: Uygun Değil (Not Confirmed), DP: Değerlendirme Yapılmadı (No Evaluation)

Birim Sorumlusu/ Department Responsible

Serkan TEKİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKGÜDAK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tasdik Olunur (Approved By)
20.10.2021

Dr. Veynel Unut ÇELEMEK
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/03/20.06.2021

Sayfa (Page) 5 / 6

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Mersis No: 26.25.0258.0550.0015

Karacaoğlu Mh. 6346 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T | +90 332 462 00 81
T | +90 332 462 00 57
E | +90 332 472 13 23



T.C.
TARIM VE ORMAN BAKANLIĞI
EUROFINS İZMİR ÖZEL GIDA KONTROL LABORATUVARI
(REPUBLIC OF TURKEY MINISTRY OF AGRICULTURE AND FORESTRY
EUROFINS İZMİR FOOD CONTROL LABORATORIES)
MUAYENE VE ANALİZ RAPORU
(ANALYSIS REPORT)

AB-0765-T
2113737
10-21

Rapor No (Report Number) : 2113737
Revizyon No (Revision No) : 00

Rapor Yayın Tarihi (Date of Issue): 03.10.2021

Analizler (Analyses)	Sonuç (Result)	ÖB (MU)	GR(%) (RM)	LOQ (LOQ)	Analiz Metodu (Analysis Method)	LDK (ML)	DS (Evaluation)
Notlar (Notes): 1. Yapılan muayene ve analiz sonuçları aşağıda belirtilen değerler toplanmıştır. (Below mentioned values have been determined from the analytical work performed.) 2. Bu analiz raporuna sadece idari işlemler ve teknik amaçlarla kullanılmalıdır. (This report shall not be used in the executive-judicial processes and for advertising purposes.) 3. Bu analiz raporunun hiçbir bölümü tek başına veya ayrı ayrı kullanılamaz. (This report with all parts is a whole, no part of this report can be used separately.) 4. Analiz sonuçları aşağıda belirtilen kapsamda geçerlidir. (Results of analysis belong to sample mentioned above.) 5. "Raporla ilgili akreditasyon kapsamıdır." ("Analysis is covered by our accreditation scope.") 6. Raporlar laboratuvarın yazılı izni olmadan başka bir kişiye verilmemelidir. (Reports may not be copied or reproduced in part without the written permission of the laboratory. Unsigned and unsealed reports are invalid.) 7. "Her operasyonun ayrı ayrı değerlendirilmesi, her yerde yerli yapılan sonuçlar ile ilgili değildir. (Except marked with *, the commented results are not associated with scope of accreditation.) 8. Laboratuvar bilgileri müşteri tarafından sağlanmıştır. Raporlarda verilen bilgilerden sorumludur. Bilgileri müşteri tarafından sağlanmıştır ve bu bilgilerin doğruluğu garanti edilemez. (The laboratory is not responsible for any deviation in the relevant results if the sample information is provided by the customer and this information may affect the validity of the results.) 9. Numune alınış laboratuvar tarafından yapılmıştır. Analiz raporunda belirtilen sonuçlar laboratuvar tarafından sağlanmıştır. (Sampling is not performed by the laboratory and sampling is the responsibility of the customer. The analysis results are valid for the sample that has been delivered to the laboratory.) 10. Kısaltmalar: (Uygun, LD Uygun Değil, DZ, Değerlendirme Yapılmadı, DS, Değerlendirme Sonucu (Abbreviations: C: Confirmed, NC: Not Confirmed, NF: No Evaluation)) 11. Türk Akreditasyon Kurumu (TAK) ile ilgili raporların tamamı, Avrupa Akreditasyon Birliği (EAB) ile Çek Tarafı Akademi ve Uluslararası Laboratuvar Akreditasyon Birliği (ILAC) ile ilgili raporların tamamı geçerlidir. (Turkish Accreditation Agency (TAK) is a signatory to the European co-operation for Accreditation (EAB) Multilateral Agreement (MLA) and to the International Laboratory Accreditation Cooperation (ILAC) Mutual Recognition Arrangement (MRA) for the recognition of test reports.) 12. Uygunluk bilgileri analiz sonucunda ilgili belirlenen sınırların dışına çıkmadığı takdirde geçerlidir. (The declaration of conformity was made according to the result obtained by subtracting the measurement uncertainty from the analysis result.) 13. Belirli edilen genişletilmiş ölçüm belirsizliği, standart belirsizliği k=2 olan genişletme katsayısı ile parçaları sonucunda 100% oranında güvenilirlik seviyesi sağlanmıştır. (The declared expanded measurement uncertainty provides a 100% confidence level as a result of multiplying the standard uncertainty by the expansion coefficient k = 2.)							

LOQ: Ölçüm Limiti (Limit of Quantification), GR: Geri Kazanım (Recovery), ÖB: Ölçüm Belirsizliği (MU: Measurement Uncertainty), LDK: Limit Değer Kaynağı (Source of Limit)
DS: Değerlendirme Sonucu (Evaluation): 10: Uygun (Confirmed), LD: Uygun Değil (Not Confirmed), DZ: Değerlendirme Yapılmadı (No Evaluation)

Birim Sorumlusu / Department Responsible

Sekban TEĞİN
Kimyasal An.Lab.Birim Sorm.
Head of Chemical Analysis Dept.

İrem AKGÜDÜK
Num.Kabul ve Raporlama Sor.
Head of Sample Admission and
Reporting Dept.

Tamam Olunmuş (Approved by)
20.10.2021

Dr. Vefail Umut ÇELİK
Laboratuvar Müdürü
Laboratory Manager

F-7.8/001/02/10.06.2021

Sayfa (Page) 6 / 6

Eurofins İzmir Gıda Analiz
Laboratuvarları Limited Şirketi
Mersis No : 0625 0250 0550 0015

Karacaöğlen Mh. 6366 Sk. No:27/A
35070 Bornova - İzmir / Türkiye
www.eurofins.com.tr

T : +90 232 462 00 81
T : +90 232 462 00 57
F : +90 232 472 13 23